

H₂O₂ SEED PRIMING TO ALLEVIATE THE SALINITY EFFECT IN TOMATO (*SOLANUM LYCOPERSICON*)

Wafa Saidi^{1,2}, Hayet Beltayef^{2,3*}, Fatma Kalleli^{1,4}, Mouna Mechri⁵, Abeer Hashem⁶, Mekhled M. Alenazi⁷, Elsayed Fathi Abd Allah⁷, Cristina Cruz⁸ and Mahmoud M. Hamdi^{1,9}

¹Department of Horticultural Sciences and Vegetable Crops, High Institute of Agronomy of ChottMariem, University of Sousse, 4042 Sousse, Tunisia

²Research laboratory: Support for sustainable agricultural productivity in the Northwest region, EL KEF Higher School of Agriculture, 7119, EL KEF, Tunisia

³Faculty of Sciences of Bizerte, University of Carthage, Tunisia

⁴LR13AGR02 Improvement and Integrated Development of Animal Productivity and Food Resources, University of Carthage, Mateur Higher School of Agriculture, 7030, Mateur, Tunisia

⁵National Institute of Field Crops, Tunisia

⁶Botany and Microbiology Department, College of Science, King Saud University, P.O. Box. 2460, Riyadh 11451, Saudi Arabia

⁷Plant Production Department, College of Food and Agricultural Sciences, King Saud University, P.O. Box. 2460, Riyadh 11451, Saudi Arabia

⁸Center for Ecology, Evolution and Environmental Changes & CHANGE-Global Change and Sustainability Institute, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal

⁹Horticultural Science Laboratory, National Agronomic Institute of Tunisia, University of Carthage, Tunisia

*Corresponding author's email address: hayet.beltayef@fsb.ucar.tn

Abstract

Salinity is one of the important constraints of sustainable agriculture, globally reducing crop production by impairing various biochemical, physiological and molecular functions. To examine hydrogen peroxide (H₂O₂) role to palliate the salinity stress, we investigated the seed-inducing effect of different H₂O₂ concentrations (0, 50, 100 and 200 mM) in tomato (*Solanumlycopersicon*) crops resistance to salt stress. During this study, we estimated the eventuality of preparing tomato seeds with H₂O₂ to promote seed sprouting, plant growth and induce salt stress tolerance. Soaked seeds were pre-germinated and grown in pots on different saline soils (0, 40 and 80 mMNaCl). Findings revealed that saltness explosively affected tomato germination and seedling growth. Seed germination with 50 and 100 mM H₂O₂ improved germination in saline solutions, while high levels of hydrogen peroxide pretreatment (200 mM) inhibited tomato germination. In the pot experiment, application of hydrogen peroxide significantly increased tomato fresh weight (85% at 200 mM dose) compared to controls. At 80 mMNaCl, the concentrations 50 and 100 mM of H₂O₂ are showing the most important effect on tomato fresh weight. Results indicate that treatments have a positive effect on membrane integrity (100% at 200 mM) and a general positive effect was observed for all properties measured under salt stress. At the same time, H₂O₂ priming significantly increased catalase activity. Seed preparation with hydrogen peroxide could be an efficient sustainable way to reduce tomato salinity stress.

Keys words: Seed priming, Hydrogen peroxide, Salt stress, Tomato, Germination, Plant growth, Plant physiology, Catalase.

Introduction

The impact of salt stress on food security has become a significant threat in the agricultural sector. The increasing salinity levels due to climate change are significantly impacting food production on new farmlands (Hussain *et al.*, 2018). In fact, soil salinity disrupts various plant physiological processes, leading to oxidative damage and alterations in leaf gas exchange, ultimately resulting in decreased plant growth (Truşcă *et al.*, 2023). High salinity levels also disrupt plant metabolism by increasing ion toxicity, reducing the availability of essential elements and affecting the synthesis of proteins and lipids (Houmani *et al.*, 2022). This leads to premature plant senescence, reduced photosynthetic efficiency, and ultimately decreased crop yields. Plants have developed mechanisms to detect external signals and respond effectively to stress. When plants encounter abiotic and biotic stressors, there is an increase in the production of reactive oxygen species (ROS), leading to oxidative damage to cellular structures

and plant molecules. Among them, we can mention the hydrogen peroxide, which a stable molecule that can freely diffuse. It plays a crucial role in voltage signaling pathways (Andrés *et al.*, 2022). It can therefore stimulate a number of adaptive responses that enhance resistance to biotic and abiotic stressors. Antioxidant systems that adjust the hydrogen peroxide levels encompass of two types of H₂O₂ scavengers: enzymatic including ascorbate peroxidase, peroxidoxin, catalase and glutathione reductase as well as non-enzymatic compounds like flavonoids, glutathione and ascorbate continuously regulate ROS concentrations in H₂O₂ (Martemucci *et al.*, 2022). Furthermore, plants have continuously developing a manner to use small amount titers of ROS as a signaling component to maintain different processes to response to environmentally stress (Tarkowski *et al.*, 2022). At higher concentrations, it contributes cell metabolites oxidative damage. But otherwise, at low concentrations it generates cell signaling (Fujii *et al.*, 2022). In particular, Redox imbalances affiliated with abiotic stress like salinity increase metabolic

rates and ultimately regulate H₂O₂ production in plant cells (Truşcă *et al.*, 2023). Over the last decade, the latest studies have revealed that pretreatment with suitable H₂O₂ amount by expressing specifics and controlling different stress-responsive pathways can enhance the abiotic stress tolerance (Sachdev *et al.*, 2021).

The small size and comparatively lengthy lifetime of H₂O₂ molecules lets them to pass via cell membranes to different cell compartment making easy various signaling function (Hasanuzzaman *et al.*, 2020). Therefore, priming is considered the greatest mechanism to induce stress tolerance in plants (Fujii *et al.*, 2022), and increasing evidence indicates that initial exposure to chemical conditioning agents makes plants more resistant to stress (Hönig *et al.*, 2023). Priming has been proved as good as a modulator for abiotic stress tolerance (Salam *et al.*, 2022). Subsequently, plants utilize an antioxidant enzymatic system to repair damage caused by FAOx. For example, superoxide dismutase (SOD) helps neutralize the O₂⁻ anion by converting it into H₂O₂, which can be harmful to the plant. Other enzymes, such as catalases (CAT) and ascorbate peroxidase (APX), are then utilized to transform H₂O₂ into water and oxygen. This process acts as a stress signal for the plant, prompting it to activate its antioxidant system (Hasanuzzaman *et al.*, 2020). Although plants able to make a spacious range of antioxidants that can mitigate oxidative damage, other exogenous applications must be further investigated, including preparing seeds with stressors or chemical mixture that make them extra proof against subsequent stress situations.

Priming is considered the greatest mechanism to induce stress tolerance in plants (Fujii *et al.*, 2022), and increasing evidence indicates that initial exposure to chemical conditioning agents makes plants more resistant to stress (Hönig *et al.*, 2023). In addition, priming has been proved as good as a modulator for abiotic stress tolerance (Salam *et al.*, 2022). This process acts as a stress signal for the plant, prompting it to activate its antioxidant system (Hasanuzzaman *et al.*, 2020). Despite priming importance from a point of view agronomic and ecologic, the fundamental mechanisms worried in processing this plant are still unknown (Tripathi *et al.*, 2024). In fact, understanding the mechanisms behind stress tolerance induced by hydrogen peroxide is valuable for identifying biotechnological approaches to improve resistance to abiotic stress. Maintaining a precise balance between H₂O₂ secretion and depletion is crucial to using H₂O₂ as a signaling molecule without causing toxicity. To evaluate the effects of H₂O₂ priming, experiments were conducted on tomato plants exposed to salinity stress. The tomato *Solanum Lycopersicon* is a highly important market garden crop globally and plays a strategic role in Tunisian agriculture, covering 13.7% of total area dedicated to market gardening. It is a popular vegetable widely consumed by both urban and rural Tunisians. However, Tunisia's heavy reliance on surface water for irrigation puts it at risk of water shortages due to changing weather patterns caused by the climate crisis (Mkaddem *et al.*, 2022). Additionally, the tomato's supply is constrained by low yields caused by various biotic and abiotic factors, including salinity. Tomatoes are known to be sensitive to salt levels and are one of the most

important vegetables worldwide (Agius *et al.*, 2022). In this study, we examined the effects of different levels of hydrogen peroxide on reducing the negative impact of soil salinity during germination. We also looked at how H₂O₂ priming can induce changes in morphology, biochemistry, and physiology to reduce the consequences of oxidative and osmotic stress caused by salinity in tomato plants.

Material and Methods

Plant material: In all the experiments, tomato seeds of the “Rio Grande” variety were used.

Seed priming with H₂O₂: Seed priming is a method that involves hydrated seeds tokickstart the germination process and activating pre-germination metabolic functions. This technique was suggested as a way to improve seed vigor before planting. During this study, tomato seeds were subjected to varying concentrations of hydrogen peroxide solutions (0, 50, 100, and 200 mM) in a dark environment at 22°C for 2 hours. They were then rinsed with distilled water to remove any excess chemicals and further cleansed with a 40% sodium hypochlorite solution for 15 minutes, followed by 3 washes with sterile distilled water.

Germination experiment: Seed germination was carried out in Murashige and Skoog Medium (Murashige & Skoog, 1962) using plastic Petri dishes. After priming period, the seeds were salinized by adding salt stress-inducing agent NaCl at various concentrations: 0, 40, 80 and 120 mM. We performed an experiment 4x4 factorial organized in randomized layout (4 salt concentrations: 0, 40, 80, and 120 Mm NaCl) and 4 concentrations of hydrogen peroxide priming (0, 50, 100 and 100 mM). Ten tomato seeds were placed in each of the 48 Petri dishes, 10 seedlings were arbitrarily selected from each Petri dish during germination period and stem length turned into measured in cm.

Germination was evaluated every 48 h and percentage germination (GP) was determined:

$$GP = \frac{\text{Number of normally germinated seeds}}{\text{Total number of seeds sown}} \times 100$$

Plant growth: An experiment was performed under greenhouse conditions. Temperature varied between 20°C on night time and 25°C on day, relative humidity was betwixt 50–70% and photoperiod fixed on 16 hours and 8 h dark. The experiment design composed of 3 nutrient solutions (control non-salt, 40 and 80 mM NaCl) and 4 H₂O₂ treatments (without application control, 50, 100 and 200 mM H₂O₂). 10 plants for one unit of experience were used for each dose of NaCl and each concentration of H₂O₂. The experimental design was set out in a randomized complete block. Globally, 3 replicates per treatment came to 36 experimental unit plots composed by 10 plants each (n = 360 plants). Before sowing, the tomato seeds (cv. Rio Grande) were soaked for 2 hours in varying doses of H₂O₂ solution. One week after seeding, seedlings were transferred into 1L plastic pots planted with unsterile substrate (soil: peat: sand (1/2: 1/4: 1/4)).

Each 3 days, plants were watered with 500 mL of water or with a solution of NaCl (40 and 80 mM).

Studied parameters

Agronomic parameters: At the final harvest, which occurred 21 days after transplanting, the fresh yield of tomatoes was evaluated in eight plants per plot. The shoot biomass was dried in a pressured-air oven at 80°C for three days to determine the shoot's dry biomass. To assess the vegetative growth of the tomato plants, the number of flowers on ten plants for each treatment and block was measured.

Physiological parameter: Membrane integrity was estimated by measuring the electrical conductivity using a digital conductivity meter. The loss of electrolytes was measured at the leaf level and it provides information on membrane integrity. Electrolyte loss was determined using the following method: 5 mm sheet slices obtained from 200 mg of fresh material are placed in tubes containing 20 ml of distilled water. These tubes are incubated at 4°C for 10 hours. We thus measure the electrical conductivity of this water, (EC1). The samples are then autoclaved at 120°C for 15 min, and then cooled to 25°C. The conductivity of the solution of the destroyed leaves, ie (EC2), is again measured. The percentage of membrane damage (Silva *et al.*, 2022) was determined as following:

$$EL = (EC1/EC2) * 100$$

Measurements were performed on 4 replicates per treatment.

Biochemical parameter: A revelation on gel of the catalase activity (CAT) was carried out on the extracts of total proteins of tomato leaves treated with different doses of Hydrogen peroxide compared to the control (Bradford, 1976).

Revelation on gel of catalase activity (CAT): The method of non-denaturing gels of polyacrylamides was prepared without sodium dodecyl sulfate (SDS). Gels are formed by a separation gel and a compression gel. The first consists of a 30% acrylamide solution of 2% bis-acrylamide, tris-HCl (pH = 8.7), H₂O, persulfate of ammonium (APS) 10% and Tetra methyl-ethylene-diamine (TEMED). Even composition for the second gel with differences in concentration and pH for the trisHCl solution that has a pH of 6.8. After polymerization of the gels, soluble proteins (50µg/sample) are loaded in the wells. The migration on gel is carried out at 70 mA and at 4°C. Finally, the step of revelation consists of incubating the gel in the dark for 30 min with shaking in a solution which contains: potassium phosphate, H₂O₂, EDTA and nitrobluetetrazolium (NBT). After rinsing with distilled water, the gel is transferred to another solution that contains potassium phosphate, H₂O₂, EDTA, acid ascorbic acid and H₂O₂. At the end the gel is incubated in the light without shaking until the appearance of catalase isomorphs (CAT) as a light band on a dark background gel (Pezzoni *et al.*, 2018).

Statistical analysis

The statistical analyses were used to compare means of different treatments, we carried out analyzes of variance and comparisons of means by the Duncan test at the 5% probability threshold using the SAS software.

Results

Germination rate: The seed germination started on the 3rd day, as well as final germination rate was 100% under the control condition (Fig. 1). Results showed no change in growth rate when 40 mMNaCl was used compared to conditions without salt. High salt concentration negatively affects tomato growth, and salt stress at 120 mMNaCl can significantly abolish this symptom. Only some seeds germinated on the 5th day, and the final germination rate declined to 40% (Fig. 2). Priming with 50 and 100 mM H₂O₂ doses enhanced the percentage germination under salt stress. In fact, treatment with 100 mM H₂O₂ appeared the most promising application in accordance with the data arranged in Figure 2. However, this positive effect of H₂O₂ on germination was limited and germination rate considerably decreased with the increased concentration of H₂O₂ (from 200 mM).




Fig. 1. Effect of NaCl salinity on germination rate of tomato seeds cv. Rio Grande.

Agronomic Parameters

Stems length: The effect of salt stress on the stem length of tomato is shown in Figures 3 and 4. Means comparison for this parameter present a highly significant effect of salt. Indeed, stem length was reduced by 70% at 120 mMNaCl (Fig. 3) in comparison with unstressed treatment which means that the salt inhibited the growth of the tomato plants. For the percentage germination, seed priming treatment with 50 mM H₂O₂ (compared to NaCl treatment) reduced salt stress and progressed the seedling overall performance (Fig. 4). As well as different effects of priming were revealed. The height of the stem is optimal for handling 50 mM H₂O₂ (9 cm) at 40 mMNaCl. However, application of 100 and 200 mM H₂O₂ contributed a decrease of stem length compared to control.




Fig. 2. Effect of different doses of H₂O₂ on germination rate of tomato seeds cv. Rio Grande at different concentrations of NaCl: a: 40 mM NaCl; b: 80 mM NaCl; c: 120 mM NaCl.




Fig. 3. Effect of different NaCl concentrations on stem length *In vitro* c.v Rio Grande tomato plants. Data are means SD of four replications, significant means at $p \leq 0.001$.




Fig. 4. Effect of four H₂O₂ concentrations applied in pretreatment of salt stress seeds (0,40, 80 and 120 mM) on stem length tomato. Data are means SD of four replications, significant means at $p \leq 0.001$.

Plant growth: A significant ($p \leq 0.05$) effect of seed priming on the fresh and dry weight of shoots grown under salt stress has been presented in Table 1. In fact, shoot fresh weight was reduced by 30.8% and 67.04% compared with the control for 40 and 80 mM NaCl. Salt treatment resulted in a decrease in shoot dry weight. This trend was more pronounced with increasing NaCl concentration. Plants salinated with 80 mM NaCl had 8.8% lower shoot dry weight than non-salinated plants (Table 1). Seed priming with H₂O₂ significantly increased shoot fresh weight in both unstressed and salt-stressed conditions. At 40 mM NaCl, priming with

200 mM H₂O₂ clearly increased shoot fresh weight by 17.44% compared with untreated plants. At 80 mM NaCl, treatment with 50 mM H₂O₂ significantly increased fresh shoot weight by 47.13% compared with untreated plants. Whereas seed priming with 200 mM H₂O₂ had no effect on shoot fresh weight compared with untreated plants at this NaCl concentration (80 mM NaCl). At 40 mM NaCl, seed priming with 100 mM H₂O₂ increased shoot dry weight in against the untreated samples. With this concentration, shoot dry weight was improved by 3.86% compared to the untreated plants. Seed priming with 200 mM H₂O₂, however, did not enhance plant dry weight. This parameter was dramatically decreased after 200 mM H₂O₂ application (58.27%).

A significant ($p \leq 0.05$) effect of different amount of Salinity on flower number. Salinity levels are to flower number. At higher salinity level, this parameter was reduced by 71.43% against control. Seed priming with 50 mM H₂O₂ raised the number of flowers by 27.26% at level 40 mM NaCl against non-primed treatment. At higher H₂O₂ level (200 mM H₂O₂), no positive effect was shown by seed priming under salinity.

Membrane integrity: Membrane integrity measurements can provide information on electrolyte loss in tomato leaf cells under salt stress, and on the effect of hydrogen peroxide in salt stress tolerance (Fig. 5). Electrolyte leakage (EL) was improved in tandem with salinity amount rising in comparison with control tomato plants. For untreated plants with H₂O₂, EL was about 30% more than in comparison to non-salinized plants. Treatments with 100 mM H₂O₂ had minimal negative impact on electrophoresis less than 120 mM NaCl amendment. Moreover, a dose of 100 mM H₂O₂ diminished 50% of damage compared to the control plants in 80 mM NaCl amendment plants. During this test, the highest leakage rates were reported in control samples tracked by 200 mM H₂O₂-primed plants stressed by 120 mM NaCl.

Catalase activity gel revealed: Figure 6 illustrated catalase activity in tomato leaves in response to seed treatment with hydrogen peroxide. Data analysis showed that seed priming strategies were accompanied by an enhancement of CAT activity. Each band corresponds to one or more isomorphs associated with four tetrameric structure views of enzyme. Our results of CAT activity revelation on gel showed that the activity level of this enzyme varied as a function of the H₂O₂ applied doses.

Table 1. Effects of hydrogen peroxide (H₂O₂) treatment on growth characteristics of tomato c.v Rio Grande under salt stress.

Parameters	NaCl (mM)	Priming treatment with H ₂ O ₂				Statistic effect
		Untreated	50 mM H ₂ O ₂	100 mM H ₂ O ₂	200 mM H ₂ O ₂	
Freshweight	0	108.65 a	95.73 b	80.14 c	70 d	***
	40	74.65 c	75 b	73.13 d	90.86 a	***
	80	3.9 c	67.9 a	53.4 b	35.5 c	***
Dry weight	0	25 a	18.10 b	17.35 b	13 c	***
	40	15.96 a	11.6 b	16.6 a	9.3 c	***
	80	22.8 a	16 c	16 c	18 b	***
Flower number	0	14 a	12 b	11 b	9 c	***
	40	8 b	11 a	9 b	8 b	***
	80	4 b	6 a	5 b	4 b	***

Data represent means \pm SD, n = 10. Different letters mean significant difference according to a Duncan's multiple range test (***) $p \leq 0.001$




Fig. 5. Effects of four concentrations of H₂O₂ pretreatment of cv. Rio Grande tomato seeds on membrane integrity under increasing salt stress conditions (0, 40, 80, 120 mM NaCl). Data are means \pm SD of four replications, significant means at $p \leq 0.001$.




Fig. 6. Non-denaturing gel reveal of catalase activity (CAT) in tomato leaves treated with three H_2O_2 concentrations (0, 50, 100, and 200 mM) compared to the control untreated plants.

Discussion

Plant growth: Plants exposed to high salinity can cause significant productivity reductions in the world which cause oxidative damage and ion toxicity (Ilangumaran & Smith, 2017). Therefore, a concerted effort to understand the effects of salinity on plants is critical to addressing the global salinity problem (Chakma *et al.*, 2019). The aim of this study was to test the effectiveness of H_2O_2 priming to induce salt tolerance in tomato plants under salt stress and to identify the main morphological, physiological and biochemical changes involved in this process.

Plant development preliminary stages are a key to anticipate plant development which is highly dependent to seedling germination (Rajabi *et al.*, 2020). This study disclosed that salinity caused a significant abatement in tomato seed germination parameters, suggesting that salt is unfavorable for tomato growth. However, these parameters vary by priming concentration (Lu *et al.*, 2022). It is well known that salinity is negatively correlated with germination (Zhao *et al.*, 2021). However, this negative correlation varied with salt concentration. At low NaCl concentrations, it induced seed dormancy whereas at high concentrations it inhibited germination due to Na^+ and Cl^- accretion (Zhao *et al.*, 2021). Our results did not reveal any impact of salt on GP at the lowest salt concentration used (40 mM). In general, salt affects germination by altering water and seed germination due to reduced osmotic pressure for germination, lower water absorption (Fasih *et al.*, 2021). Furthermore, high concentrations of Na^+ ions in the medium can lead to osmotic stress resulting in decreased water uptake by plant tissues (Polash *et al.*, 2019). In addition, salt could modify intoxication due to ion toxicity. Disruption of enzymatic activity leads to significant changes in plant growth, such as changes in nucleic acid and protein transfer (Peng *et al.*, 2023), disruption of hormone abundance and reduced reserves (Sachdev *et al.*, 2021). Additionally, salt can stimulate phenolic compounds, which can decrease germination rate (Peng *et al.*, 2023). But also other factors can influence seed germination under saline-alkaline conditions like water, temperature and light (Wang *et al.*, 2022). For shoot and root lengths, this parameter decreased with increasing salt concentration. In fact, roots are the first ones that come into direct contact with salinity soil. They soak up the water and deliver it to aerenchyma organs (Wang *et al.*,

2022). Indeed, plant tissues can be severely affected by salinity. The decrease in stem length of tomato seedlings observed in this study with increasing salinity elucidated that salinity reduces the intercellular carbon dioxide concentration, which in turn reduces the rate of photosynthesis through stomatal closure. Under high salt levels, Na^+ can cause lower transport rate of essential ions such as NO_3^- that reduce the N-containing compounds and ultimately inhibit plant growth (Khaleduzzaman *et al.*, 2021). In fact, H_2O_2 has a positive effect on the response of plants to different types of stress (Danial & Basset, 2024). The results of this study showed that plants treated with H_2O_2 showed a higher germination rate compared to plants treated with NaCl. The optimal concentration of H_2O_2 for germination was 100 mM and 50 mM for Stem length. Seed priming with H_2O_2 significantly increased germination in comparison with the control which is confirmed by results of several reports (Migahid *et al.*, 2019; Marthandan *et al.*, 2020). They have revealed a positive effect of priming on plant growth parameters. Early initiation of transplantation and the maintenance of decomposition and accumulation may account for the rapid germination of the first seeds. The reasons for the above-mentioned effects of seed preparation treatments may be different. Since H_2O_2 is considered as one of the signaling molecules, they may reprogram gene expression (Danial & Basset, 2024), resulting in de novo protein synthesis resulting in a membrane repair mechanism and more storage proteins and other substrates to enhance and synchronize germination. In plants, H_2O_2 is widely used in various biochemical and physiological activities. Ergo its long lifespan and flexibility, H_2O_2 can cross cell membranes and can act as a signaling molecule in stress signaling pathways. These processes can produce different stress-related responses. This research showed that seed germination with an optimal amount of H_2O_2 can enhance plant growth under salinity conditions. H_2O_2 as the crucial agent revealed a rising in dry weight of cannabis crop (Islam *et al.*, 2022). Furthermore, seeds grown with H_2O_2 exhibited exorbitant bigger dry weight and root length than wheat under drought conditions (Habib *et al.*, 2020). It has been suggested that seed germination leaves a "stress memory" that activates stress response mechanisms after germination (Lutts *et al.*, 2016).

However, seed germination and increased H_2O_2 decreased the number of sprouts. Despite acting as a signaling molecule, excess H_2O_2 induces oxidative stress, which damages cells and leads to systemic death (Habib *et al.*, 2020). Number of flowers was also bettered by H_2O_2 priming. Priming with 50 mM H_2O_2 was the most effective for perfecting number of flowers. It has been conjectured that seed priming may trigger stress responsive set up corresponding to antioxidant stimulation, abscisic acid signaling, and hormonal modulation (Rhaman *et al.*, 2020). It has been suggested that seed priming system can engender physiological acclimatization, that lead to its rise crop yield under stress situation that lead to its increase (Marthandan *et al.*, 2020). In fact, similar results obtained for wheat yield which was raised under drought condition after priming with suitable amount H_2O_2 (Marthandan *et al.*, 2020). Appropriate amount of H_2O_2 can induce stress tolerance by regulating multiple physiological processes such as photosynthesis, stomatal movement, uptake adaptation, and ROS detoxification (Habib *et al.*, 2020).

Membrane integrity: Salt tolerance depended to membrane integrity mechanism. The plasma membrane is a major site of ion-specific salt damage (Sayed&Gadallah, 2019). For that, EL of the plasma membrane is the way to identify halotolerant plants. The results obtained for this parameter prove that the values of electrolyte leakage are directly proportional to increasing salt content. Likewise, the indices of lettuce leaves and roots gradually increased with rising NaCl concentration (Kazamel *et al.*, 2024). Some authors have exhibited this finding in cucumber (Soufi *et al.*, 2023), and strawberry (Wang *et al.*, 2019). In our study, the results of priming seeds with H₂O₂ reduced ion leakage in salinity-stressed tomato plants. This may be due to the stimulation of plasma membrane activity under salt stress by Hydrogen peroxide (Forghani *et al.*, 2021). In the same context, spraying sunflower leaves with H₂O₂ stimulates salt tolerance. On the other hand, in untreated plants, an overproduction of free energy with an excessive secretion of ROS contributes to an aggravated electrolyte leakage (Lephatsi *et al.*, 2021). On the other hand, treatment of plants with H₂O₂ reduced the negative effects of NaCl in a similar way to control plants. The immune system manages so-called antioxidant enzymes with the capacity to trap ROS, such as catalase (CAT). CAT is involved in the cleavage of H₂O₂ into water and oxygen. In fact, the increase in antioxidant activity by H₂O₂ priming is reflected by the stimulation of the expression of various antioxidant enzyme genes, such as superoxide dismutase, catalase and others (Lephatsi *et al.*, 2021; Savvides *et al.*, 2016). Indeed, overexpression of these enzymes can correct the negative effect of salt stress by helping to maintain redox homeostasis. Citing Example, catalase activity is significantly increased in leaves and roots in response to H₂O₂ priming, playing a key role in ROS detoxification (Kesawat *et al.*, 2023).

Conclusion

In conclusion, priming seeds with H₂O₂ contributed to ameliorate germination and growth traits of tomato under salt stress. 100 mM of H₂O₂, used as the initiator, was the most effective concentration as indicated by percentage germination and plant biomass. Seed priming was accompanied by an improvement in activity and a decrease in the amount of ion leakage. Antioxidant signaling appears to be crucial targets of H₂O₂ that lead to increased ion osmotic stress tolerance. It appears that an appropriate concentration of H₂O₂ can be used to mitigate the bad effects of salinity without influencing plants processes to grow under conditions without stress. Limits in NaCl tolerance following H₂O₂ treatment were observed during our trials. And at high concentrations, H₂O₂ can amplify the harmful effects of NaCl. Further clarify the role of H₂O₂ in salt stress tolerance by increasing the duration of soaking using the concentrations that gave the best results. Understand more about the relationship between hydrogen peroxide and the CAT enzyme. And exploit it as an early selection criterion in improvement and creation new varieties tolerant to salt stress.

Acknowledgement

The authors would like to extend their sincere appreciation to the Researchers Supporting Project Number (RSP2024R134), King Saud University, Riyadh, Saudi Arabia.

References

- Agius, C., S. von Tucher and W. Rozhon. 2022. The effect of salinity on fruit quality and yield of cherry tomatoes. *Horticulturae*, 8(1): 59.
- Andrés, C.M.C., J.M. Pérez de la Lastra, C.A. Juan, F.J. Plou and E. Pérez-Lebeña. 2022. Chemistry of hydrogen peroxide formation and elimination in mammalian cells, and its role in various pathologies. *Stresses*, 2: 256-274.
- Bradford, M.M. 1976. A Rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-Dye Binding. *Anal. Biochem.*, 72: 248-254.
- Chakma, P., M.M. Hossain and M.G. Rabbani. 2019. Effects of salinity stress on seed germination and seedling growth of tomato. *JBAU.*, 17(4): 490-499.
- Danial, A.W. and R.A Basset. 2024. Amelioration of NaCl stress on germination, growth, and nitrogen fixation of *Viciafaba* at isosmotic Na-Ca combinations and *Rhizobium*. *Planta*, 259(3): 69.
- Fasih, U.H., L. Cai, A.C. Jeffrey, A.C Sardar, W. Jun, Z. Renzhi, W. Ma and F. Muhammad. 2021. Cadmium toxicity in plants: Impacts and remediation strategies, Ecotoxicology and Environmental Safety. *Ecotoxicol. Environ. Saf.*, 211: 111887.
- Forghani, A., A.H. Forghani, M. Altafi, K. HashemiMajd and O. Sofalian. 2021. The effects of different sources of potassium and calcium on yield and ionic balance of tomatoes under salinity stress in hydroponic cultivation. *NBR.*, 8(3): 206-219.
- Fujii, J., T. Homma and T. Osaki. 2022. Superoxide radicals in the execution of cell death. *Antioxidants*, 11: 501.
- Habib, N., Q. Ali, S. Ali, M.T. Javed, M.Z. Haider, R. Perveen, M.R. Shahid, M. Rizwan, M.M. Abdel-Daim, A. Elkelish and M. Bin-Jumah. 2020. Use of nitric oxide and hydrogen peroxide for better yield of wheat (*Triticum aestivum* L.) under water deficit conditions: growth, osmoregulation, and antioxidative defense mechanism. *Plants*, 9: 285.
- Hasanuzzaman, M., M.H.M.B. Bhuyan, F. Zulfiqar, A. Raza, S.M. Mohsin, J.A. Mahmud, M. Fujita and V. Fotopoulos. 2020. Reactive oxygen species and antioxidant defense in plants under abiotic stress: Revisiting the crucial role of a universal defense regulator. *Antioxidants*, 9: 681.
- Hönig, M., V.M. Roeber, T. Schmülling and A. Cortleven. 2023. Chemical priming of plant defense responses to pathogen attacks. *Front. Plant Sci.*, 14:1146577.
- Houmani, H., J.M. Palma and F.J. Corpas. 2022. High salinity stimulates the adaptive response to potassium deficiency through the antioxidant and the NADPH-generating systems in the roots and leaves of the halophyte *cakilemaritima*. *J. Plant Growth Regul.*, 42: 6286-6306.
- Hussain, M., S. Farooq, W. Hasan, S. Ul-Allah, M. Tanveer, M. Farooq and A. Nawaz. 2018. Drought stress in sunflower: Physiological effects and its management through breeding and agronomic alternatives. *Agri. Water. Manag.*, 201: 152-166.
- Ilangumaran, G. and D.L. Smith. 2017. Plant growth promoting rhizobacteria in amelioration of salinity stress: A systems biology perspective. *Front. Plant Sci.*, 23(8):1768.
- Islam, M.J, B.R. Ryu, M.H. Rahman, M.S Rana, E.J. Cheong, M.H. Wang, J.D. Lim, M.A. Hossain and Y.S. Lim. 2022. Cannabinoid accumulation in hemp depends on ROS

- generation and interlinked with morpho-physiological acclimation and plasticity under indoor LED environment. *Front. Plant Sci.*, 5(13): 984410.
- Kazamel, A.M., S.A. Haroun, A.A. Noureldin, G.A. Sherbiny, O.A. El-Shahaby, M.R. Sofy, A.F. AlBakry and R.M.E. Gamel. 2024. Ultrastructural, secondary metabolite, and antioxidant modulation in response to salt-affected habitats induced oxidative stress and their accumulation in *Malvaparviflora* L. and *Rumexdentatus* L. *J. Soil Sci. Plant Nutr.*, 24: 389-407.
- Kesawat, M.S., N. Satheesh, B.S. Kherawat, A. Kumar, H.U. Kim, S.M. Chung and M. Kumar. 2023. Regulation of reactive oxygen species during salt stress in plants and their crosstalk with other signaling molecules-current perspectives and future directions. *Plants*, 12: 864.
- Khaleduzzaman, M., M.A. Hossain, M.J.H. Bhuiyan, S. Mahmud, M. Tahjib-Ul-Arif and Y. Murata. 2021. Chitosan mitigates salt stress in rice by enhancing antioxidant defense system. *Fundam. Appl. Agric.*, 6(4): 336-348.
- Lephatsi, M.M., V. Meyer, L.A. Piater, I.A. Dubery and F. Tugizimana. 2021. Plant responses to abiotic stresses and rhizobacterialbiostimulants: metabolomics and epigenetics perspectives. *Metabolites*, 11(7): 457.
- Lu., Y, H. Liu, Y. Chen, L. Zhang, K. Kudus and J. Song. 2022. Effects of drought and salt stress on seed germination of ephemeral plants in desert of northwest China. *Front. Ecol. Evol.*, 10: 1026095.
- Lutts, S., P. Benincasa, L. Wojtyla, S.S. Kubala, R. Pace, K. Lechowska, M. Quinet and M. Garnczarska. 2016. Seed priming: new comprehensive approaches for an old empirical technique. In: (Ed.): Araújo, S. New Challenges in Seed Biology-Basic and Translational Research Driving Seed Technology. Rijeka, *Intech. Open.*, 1-47.
- Martemucci, G., C. Costagliola, M. Mariano, L. D'andrea, P. Napolitano and A.G. D'Alessandro. 2022. Free radical properties, source and targets, antioxidant consumption and health. *Oxygen*, 2: 48-78.
- Marthandan, V., R.K. Greetha, V.G. Kumutha, A. Ranganathan, A. Karthikeyan and J. Ramalingam. 2020. Seed priming: A feasible strategy to enhance drought tolerance in crop plants. *Int. J. Mol. Sci.*, 21: 8258.
- Migahid, M.M., R.M. Elghobashy, L.M. Bidak and A.W. Amin. 2019. Priming of *Silybum marianum* (L.) Gaertn seeds with H₂O₂ and magnetic field ameliorates seawater stress. *Heliyon*, 5(6): e01886.
- Mkaddem, C. and S. Mahjoubi. 2022. Climate change and its impact on water consumption in Tunisia: Evidence from ARDL approach. Tunis, Tunisia: University of Tunis.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Physiol. Plant*, 15: 473-497.
- Peng, M., Z. Jiang, F. Zhou and Z. Wang. 2023. From salty to thriving: plant growth promoting bacteria as nature's allies in overcoming salinity stress in plants. *Front. Microbiol.*, 23(14): 1169809.
- Pezzoni, M., R.A. Pizarro and C.S. 2018. Costa. detection of catalase activity by polyacrylamide gel electrophoresis (PAGE) in cell extracts from *Pseudomonas aeruginosa*. *Biol. Protoc.*, 5: 8(11): e2869.
- Polash, MAS., MA. Sakil and M.A. Hossain. 2019. Plants responses and their physiological and biochemical defense mechanisms against salinity: A review. *Trop. Plant Res.*, 6(2): 250-274.
- Rajabi, D.A., M. Zahedi, A. Ludwiczak, S.C. Perez and A. Piernik. 2020. Effect of salinity on seed germination and seedling development of sorghum (*Sorghum bicolor* (L.) Moench) genotypes. *Agronomy*, 10(6): 859.
- Rhaman, M.S., S. Imran, F. Rauf, M. Khatun, C.C. Baskin, Y. Murata and M. Hasanuzzaman. 2020. Seed priming with phytohormones: an effective approach for the mitigation of abiotic stress. *Plants*, 25: 10(1):37.
- Sachdev, S., S.A. Ansari, M.I. Ansari, M. Fujita and M. Hasanuzzaman. 2021. Abiotic stress and reactive oxygen species: Generation, signaling, and defense mechanisms. *Antioxidants*, 10: 277.
- Salam, A., M.S. Afridi, M.A. Javed, A. Saleem, A. Hafeez, A. Khan, M. Zeeshan, B. Ali, W. Azhar and W. Sumaira. 2022. Nano-Priming against abiotic stress: A way forward towards sustainable agriculture. *Sustainability*, 14(22): 14880.
- Savvides, A., S. Ali, M. Tester and V. Fotopoulos. 2016. Chemical priming of plants against multiple abiotic 271 stresses: mission possible? *Trends. Plant Sci.*, 21: 329-340.
- Sayed, S. and M. Gadallah. 2019. Hydrogen peroxide supplementation alleviates the deleterious effects of cadmium on photosynthetic pigments and oxidative stress and improves growth, yield and pods quality of pea (*Pisum sativum* L.) plants. *Acta Physiol.*, 41: 113.
- Silva, L.D.A., J.A.G. Gomes, P.A. Rodrigues, R.P. Araújo, L.S. Pereira, A.C. Dalmolin and J.F. Amaral. 2022. Antioxidative and osmorregulatory responses of *Jatropha curcas* to water stress are genotype dependent. *AJCS.*, 16(02): 192-200.
- Soufi, H.R., H.R. Roosta, F. Fatehi and M. Ghorbanpour. 2023. Spectral composition of LED light differentially affects biomass, photosynthesis, nutrient profile, and foliar nitrate accumulation of lettuce grown under various replacement methods of nutrient solution. *Food Sci. Nutr.*, 10: 11(12): 8143-8162.
- Tarkowski, Ł.P., S. Signorelli, M.J. Considine and F. Montrichard. 2022. Integration of reactive oxygen species and nutrient signalling to shape root system architecture. *Plant Cell Environ.*, 46(2): 379-390.
- Tripathi, D. K., J. A. Bhat, C. Antoniou, N. Kandhol, V.P. Singh, A.R. Fernie and V. Fotopoulos. 2024. Redox regulation by priming agents towards a sustainable agriculture. *Plant and Cell Physiology*, pcae031.
- Truşcă, M., Ş. Gâdea, R. Vidican, V. Stoian, A. Vâtcă, C. Balint, V.A. Stoian, M. Horvat and S. Vâtcă. 2023. Exploring the research challenges and perspectives in ecophysiology of plants affected by salinity stress. *Agriculture*, 13: 734.
- Wang, J.Z., C. Jin, Y.P. Wang and B.Q. Chen. 2019. Effects of salt stress on antioxidant system activity and peroxidation damage in root tip cells of strawberry. *AJB.*, 18(28): 702-706.
- Wang, W., F. Zhang, L. Sun, L. Yang, Y. Yang, Y. Wang, K.H.M. Siddique and J. Pang. 2022. Alkaline salt inhibits seed germination and seedling growth of canola more than neutral salt. *Front. Plant Sci.*, 27(13): 814755.
- Zhao, S., Q. Zhang, M. Liu, H. Zhou, C. Ma and P. Wang. 2021. Regulation of plant responses to salt stress. *Int. J. Mol. Sci.*, 22(9): 4609.