

RESPONSE OF WHEAT GROWTH AND YIELD TO VARIOUS LEVELS OF COMPOST AND ORGANIC MANURE

MUHAMMAD IBRAHIM^{1*}, ANWAR-UL-HASSAN¹, MUHAMMAD IQBAL¹
AND EHSAN ELAHI VALEEM²

¹*Inst. Soil & Environ. Sci., University of Agriculture, Faisalabad-38040, Pakistan.*

²*Public Private Partnership Unit (PPP Unit), Planning and Development Department,
Government of Sindh, Karachi-74200, Pakistan.*

Abstract

Organic manures are traditionally used for supplying plant nutrients. Their volume and other operational problems forced the farmers to use inorganic fertilizers. The recycling of organic waste is considered one of the major solutions to get rid of huge piles of wastes. The product obtained provides nutrients in an efficient way along with improving the soil conditions. Because the current practice of using inorganic fertilizers may not fulfil and maintain the soil quality needed for sustainable crop productivity. The impact of organic manure and compost on productivity of wheat (*Triticum aestivum* L. cv. Inqlab-91) was investigated in sandy clay loam soil. The amounts of various organic manures to supplement the inorganic fertilizers must be optimized to increase crop yield. Changes in growth and yield parameters of wheat relative to inorganically fertilized controls were measured. Organic amendments had positive but variable effects. The organic manures application increased the wheat yield by 11.13 (105 %) to 13.53 (128 %) g pot⁻¹, relative to the control. The wheat plant height, number of tillers, spike length, straw yield, grain yield and 1000-grain weight all were statistically different from that of control. The findings of the trial suggested that crop productivity may be improved significantly by the application of various organic manures for longer time. Hence, instead of using inorganic chemical fertilizer alone, the integrated use could be more effective and sustainable for environment and agriculture.

Introduction

The organic matter is used since times to improve soil health and supplying plant nutrients. Various types and sources of organic wastes are utilized in agriculture but most of these materials remain unutilized, especially in resource poor countries. These practices pose serious threat to the environment and loss of useful pool of plant nutrients. The organic materials are available in bulk amounts as farm manure, city waste, poultry manure and wastes from industry like food, sugar, cotton and rice. If these materials are accumulated, these may become a potential source of air, land and water pollution. According to Economic Survey (Anon., 2007) no proper waste collection and disposal system is in place in Pakistan. The farmers are using various types of waste materials without knowing their economic benefit.

Huge amounts of organic materials are in the form of farm waste, city waste, sewage sludge, poultry litter and agro-industrial wastes (Lal, 2005; Kolay, 2000). Less than 60 % waste is collected in urban areas of Pakistan because there is new system for city waste collection and disposal (Anon., 2006). The various forms of organic manures are also being used in the fields for sustainable crop yield. But due to varying composition and water level and bulky nature, their application and transportation is a very big question in this fuel conscious world. The application also implies certain labour requirements, which add to the input cost and that is very un-economical for a resource poor farmer.

*Corresponding authors: ebrahem.m@gmail.com

Composting provides an effective and environment friendly procedure of organic waste disposal (Millner *et al.*, 1998) because it is more economical and environment friendly. It also conserves natural resources and improves cycling of non-renewable resources. Keeping in view the present energy crises it is an excellent option for energy conservation because a lot of energy is utilized in fertilizer sector. This process biologically converts the organic waste material into stable humus like substances, which may be stored and applied without any environmental impacts (Gallardo-Larva & Nogales, 1987). The organic manures and compost are important in sustaining farming by providing plant N-supply (Korsaeth *et al.*, 2002).

The compost production for agriculture has got considerable attention in socio-economic conditions in Pakistan due to sudden increase in fertilizer prices and increased depletion of soil quality. This might be due to increased costs for solid waste management and increased public awareness for wastes recycling. The process of composting provides sound way to manage large volumes of organic wastes in a comprehensive manner (Lasaridi & Stetiford, 1999). The organic matter in various forms and at various stages of decomposition has been used in soil in tons ha^{-1} for improvement of crop productivity (Terrance *et al.*, 2004). The use of inorganic chemical fertilizers is essential for crop nutrition in order to maximize productivity. The crop yields especially of wheat are stagnant for the last couple of years. The use of organic manures and composted organic materials along with chemical fertilizers may be effective for further increase in crop yield.

Pakistan is facing a severe challenge of acute food and energy crisis and sufficient food production at least for ever growing population at 2.6 percent per annum (Qazilbash, 2002) should be made possible by the use of all available resources. It seems to be very quick and efficient method to use the mineral fertilizers but their energy costs are too high and that might not be environmental friendly and constraints may be in their use by the resource poor farmers. There are very little options available for efficient food production through the available poor resources. Attention should be focused on the efficient utilization of resources by the cereals especially wheat that is of supreme importance in Pakistani agriculture due to its cultivated area and production (Anon., 2007). For these investigations such survey should be carried out so that a comprehensive technology and production package may be developed for efficient utilization of all on farm available resources. Some work in this regard has been done by Ahmad *et al.* (2008); Zahir *et al.* (2007) and Sial *et al.* (2007). The present investigations were, therefore, performed to compare raw organic manure, compost and inorganic fertilizer for wheat growth and yield.

Materials and Methods

Pot studies were conducted in the net house to see the effectiveness of raw organic manure and compost and for improving the growth and yield of wheat. Both raw organic manure and compost materials were analyzed (Fig. 1) for carbon content (Nelson & Sommers, 1996), and macro-nutrients (Ryan *et al.*, 2001). The soil was collected, air-dried, ground, sieved (2 mm mesh) and analyzed for physico-chemical properties. The soil used for this experiment was sandy clay loam (53, 20.5 and 26.5 % sand, silt and clay, respectively) having a pH of 7.54; EC_e , 1.42 dS m^{-1} , saturation percentage of 33 %. It also contained organic matter 0.59 %; total N 0.046%, available P 7.5 mg kg^{-1} and available K 121.7 mg kg^{-1} soil.

Details of the treatments given are described below:

T_1 = Control

T_2 = Organic manure at 10 Mg ha^{-1}

T_3 = Organic manure at 20 Mg ha^{-1}

T_4 = Organic manure at 30 Mg ha^{-1}


T_5 = Organic manure at 40 Mg ha^{-1}

T_6 = Compost at 300 kg ha^{-1}

T_7 = Compost at 400 kg ha^{-1}

T_8 = Compost at 500 kg ha^{-1}

The pots were filled with sieved soil (15 kg pot^{-1}). In control, recommended N, P and K fertilizers at 120 , 100 and 60 kg ha^{-1} , respectively were mixed as urea, single super phosphate (SSP) and sulphate of potash (SOP). The organic manure and composts were applied according to the treatment by mixing it with soil before filling of pots.



The values of C, N, P and K are in % while C/N, C/P and C/K are ratios.

Fig. 1. Analysis of organic manure and compost used in the experiment.

Eight seeds of wheat cultivar Inqlab-91 were sown in each pot and thinned to four plants. Canal water was used for irrigation throughout the experimental period. The pots were arranged according to completely randomized design with three replications. The wheat plants were harvested at maturity and data regarding growth and yield parameters were recorded. The data thus obtained were subjected to statistical analysis (Steel & Torrie, 1980). The treatment means were compared by least significant difference (LSD) test.

Results and Discussion

The organic manure and compost significantly increased the plant height, number of tillers, spike length, straw/grain yield and 1000-grain weight over untreated control (Table 1). The Maximum increase (16 % over control) in plant height was recorded by the application of compost at 500 kg ha⁻¹. The organic manure at 10 Mg ha⁻¹ also increased the plant height (44.06 cm). The maximum number of tiller pot⁻¹ (29) was also recorded in T₈ and T₆ where compost was applied at 500 and 300 kg ha⁻¹. The spike length was non-significant and the maximum was observed in T₈ and the minimum was recorded in T₂, T₃ and T₇. Dixit & Gupta (2000); Selvakumari *et al.*, (2000) and Khoshgoftarmanesh & Kalbasi (2002) had also concluded that crop growth may be improved by the use of organic materials.

Wheat gave highly significant response to organic manure and compost application. A comparison of fresh weight with that of control, revealed an increase of 44 % at T₇ and 27 % at T₄ (Table 1). The experimental soil contained less amount of organic matter and had a low N₂ supplying capacity. The response of wheat to fertilizer and manures on such soils had been reported to control wheat yields (Sarwar *et al.*, 2007, 2008). Addition of various levels of organic manure and compost to wheat compensated for chemical fertilizer, which might give a substitution under field conditions (Rekhi *et al.*, 2000). The wheat grown gave a significant grain yield of 14.08, 13.98 and 13.53 g pot⁻¹ in treatments where compost at 400 kg ha⁻¹, compost at 500 kg ha⁻¹ and organic manure at 30 Mg ha⁻¹ was applied. This resulted 33, 32 and 28 % more grains over control (Table 1). Previous studies have shown that organic materials (compost, manures) enhance nutrient use efficiency by slow releasing of nutrients and reducing their losses (Chang & Janzen, 1996; Paul & Clark, 1996; Muneshwar *et al.*, 2001; Nevens & Reheul, 2003). The addition of organic fertilizer increases phosphorus mobilization and soil microbial activities; it might also contribute in improving nutrition as well as crop root system.

The data regarding fresh weight of plants per pot also showed significant results at varying levels of organic manure and compost. It is evident from the data that the maximum fresh weight (57.91 g pot⁻¹) was recorded in T₇ followed by T₄. The increase in fresh weight has also been reported by Sarwar *et al.* (2008) concluding that favourable environment results in an increased activity and increased water and nutrient use efficiency resulting in more succulent plants (Ahmad *et al.*, 2008; Sial *et al.*, 2007).

Total oven dry biomass of wheat increased significantly over control with the use of organic manure and compost. The minimum oven dry weight (19.57 g pot⁻¹) was recorded in T₆ which was at par with T₈ and T₃. The reason for this increase might be the efficient use of all available resources for plant and roots because of low and continued supply of nutrients as well as more water absorption (Jagadeeswari & Kumaraswamy, 2000; Swarup & Yaduvanshi, 2000). The increase in maize growth with the use of organic materials has also been observed by Muhammad *et al.* (2007) and Iqbal *et al.*, (2008). As for as 1000-grain weight is concerned, the produce at various levels of organic manure and compost revealed that the maximum 1000-grain weight (45.26 g) was produced when organic manure was applied at 30 Mg ha⁻¹. The difference in grain yield among different levels of manure and compost was also significant. The difference in nutrient absorption greatly influences growth and yield potential (Ahmad *et al.*, 2008). These findings are in support of previous findings of Bajpai *et al.*, (2002) and Pooran *et al.*, (2002) who concluded that compost application improved all the growth parameters.

Table 1. Effect of organic manure and compost on growth and yield of wheat.

Treatments	Plant height (cm)	No. of tillers pot ⁻¹	Spike length (cm)	Spikelets per spike	Fresh weight (g pot ⁻¹)	Oven dry weight (g pot ⁻¹)	Grain yield (g pot ⁻¹)	1000-grain weight
T ₁ = Control	41.47	24.7	8.6	13.2	40.18	15.37	10.54	40.32
T ₂ = OM at 10 Mg ha ⁻¹	44.06	26.0	8.5	13.0	46.85	16.49	11.13	37.93
T ₃ = OM at 20 Mg ha ⁻¹	40.06	21.7	8.5	13.9	40.89	18.26	11.82	40.42
T ₄ = OM at 30 Mg ha ⁻¹	42.38	24.0	8.8	13.7	51.18	17.11	13.53	45.26
T ₅ = OM at 40 Mg ha ⁻¹	43.77	25.0	8.6	12.9	36.25	16.59	12.96	43.79
T ₆ = Compost at 300 kg ha ⁻¹	41.90	29.0	8.9	13.7	45.56	19.57	13.66	42.76
T ₇ = Compost at 400 kg ha ⁻¹	40.47	28.3	8.5	13.9	57.91	17.80	14.08	42.39
T ₈ = Compost at 500 kg ha ⁻¹	48.11	29.0	9.0	14.4	34.16	19.18	13.98	44.03
LSD _{0.05}	5.29	5.43	0.88	1.08	16.90	3.52	2.82	3.99

OM = Organic Manure

The present investigations have demonstrated the improvement of wheat growth and yield with the use of organic manure and compost when they were compared with chemical fertilizer. It is quite possible to get higher wheat yield by the integrated use of organic and inorganic fertilizers. This will result in soil quality improvement, which should be investigated under field conditions.

Acknowledgements

The authors are very obliged for extensive work and criticism by Late Dr. Muhammad Younis on the earlier draft of the manuscript. The work presented here is a part of Ph.D. thesis.

References

Ahmad, R., M. Naveed, M. Aslam, Z.A. Zahir and M. Arshad. 2008. Economizing the use of nitrogen fertilizer in wheat production through enriched compost. *Renew. Agric. & Food System.*, 23(3): (In press).

Anonymous. 2006. Economic Survey of Pakistan 2004-2005. Government of Pakistan, Finance Division, Economic Advisory Wing, Islamabad.

Anonymous. 2007. Economic Survey of Pakistan 2006-2007. Government of Pakistan, Finance Division, Economic Advisory Wing, Islamabad,

Bajpai, R.K., S.K. Upadhyay, B.S. Joshi and R.S. Tripathi. 2002. Productivity and economics of rice (*Oryza sativa L.*)-wheat (*Triticum aestivum L.*) cropping system under integrated nutrient supply systems. *Indian J. Agron.*, 47: 20-25.

Chang, C. and H.H. Janzen. 1996. Long-term fate of nitrogen from annual feedlot manure application. *J. Environ. Qual.*, 25: 785-790.

Dixit, K.G. and B.R. Gupta. 2000. Effect of Farmyard manure, chemical and Biofertilizers on yield and quality of rice (*Oryza sativa* L.) and soil properties. *J. Indian Soc. Soil Sci.*, 48: 773-780.

Gallardo-Larva, F. and R. Nogales. 1987. Effect of application of town refuse compost on the soil plant system – A review. *Biol. Wastes.*, 19: 35-62.

Iqbal, M., A. Hassan and M. Ibrahim. 2008. Effects of tillage systems and mulch on soil physical quality parameters and maize (*Zea mays* L.) yield in semi-arid Pakistan. *Biol. Agric. & Hort.*, 25(4): 311-325.

Jagadeeswari, P.V. and K. Kumaraswamy. 2000. Long-term effects of manure-fertilizer schedules on the yield of and nutrient uptake by rice crop in a permanent manorial experiment. *J. Indian Soc. Soil Sci.*, 48: 833-836.

Khoshgoftarmash, A.H. and M. Kalbasi. 2002. Effect of municipal waste leachate on soil properties and growth and yield of rice. *Commun. Soil Sci. Plant Analysis.*, 33 (13&14): 2011-2020.

Kolay, A.K. 2000. Basic Concepts of Soil Science. 2nd Ed. New-Age International Publisher, New Delhi. 256 p.

Korsaeth, A., T.M. Henriksen and L.R. Bakken. 2002. Temporal changes in mineralization and immobilisation of N during degradation of plant material: implications for the plant N supply and nitrogen losses. *Soil Biol. Biochem.*, 34: 789-799.

Lal, R. 2005. World crop residues production and implications of its use as a bio-fuel. *Environ. Intl.*, 31: 575-584.

Lasaridi, K.E. and E.I. Stentiford. 1999. Composting of source separated MSW: an approach to respirometric techniques and biodegradation kinetics. In: *International Symposium on "Composting of Organic Matter"*. 31-September, 1999, Kassandra-Chalkidiki, Greece.

Millner, P.D., L.J. Sikora, D.D. Kaufman and M.E. Simpson. 1998. Agricultural uses of biosolids and other recyclable municipal residues. p. 9-44. In: *Agricultural Uses of Municipal, Animal and Industrial Byproducts*. Conservation Research Reports 44. R.J. Wright, W.D. Kemper, P.D. Millner, J.F. Power and R.F. Korcack (eds.). USDA Agricultural Research Survey, Washington, DC.

Muhammad, S., T. Muller and R.G. Joergensen. 2007. Compost and P amendments for stimulating micro-organisms and maize growth in a saline soil from Pakistan in comparison with a non saline soil from Germany. *J. Plant Nutr. & Soil Sci.*, 170 (6): 745-751.

Muneshwar, S., V.P. Singh, K.S. Reddy and M. Singh. 2001. Effect of integrated use of fertilizer nitrogen and farmyard manure or green manure on transformation of N, K and S and productivity of rice-wheat system on a Vertisol. *J. Indian Soc. Soil Sci.*, 49: 430-435.

Nelson, D.W. and L.E. Sommers. 1996. Total carbon, organic carbon and organic matter. p. 961-1010. In: *Methods of Soil Analysis*: Part 3-Chemical Methods. (Ed.): J.M. Bigham. American Soc. Agron. Inc. Madison, Wisconsin.

Nevens, F. and D. Reheul. 2003. The application of vegetable, fruit and garden waste (VFG) compost in addition to cattle slurry in a silage maize monoculture: nitrogen availability and use. *Europ. J. Agron.*, 19: 189-203.

Paul, E.A. and F.E. Clark. 1996. *Soil microbiology and biochemistry*. Academics Press, San Diego, CA.

Pooran, C., P.K. Singh, M. Govardhan and P. Chand. 2002. Integrated management in rainfed castor (*Ricinus communis*). *Indian Prog. Agric.*, 2: 122-124.

Qazilbash, A.A. 2002. Population growth and its impact on environment. *Population & E Bulletin*, Vol. 2: 4.

Rekhi, R.S., D.K. Benbi and B. Singh. 2000. Effect of fertilizers and organic manures on crop yields and soil properties in rice-wheat cropping system. Page 1-6 In: *Long-term soil fertility experiments in rice-wheat cropping systems*. (Eds.): Abrol, I.P., K.F. Bronson, J.M. Duxbury and R.K. Gupta. Rice-Wheat Consortium Paper Series 6. New Delhi, India: Rice-Wheat Consortium for the Indo-Gangetic Plains.

Ryan, J., G. Estefan and A. Rashid. 2001. Soil and Plant Analysis: Laboratory Manual. International Centre for Agricultural Research in Dry Areas (ICARDA), Aleppo. 172p.

Sarwar, G., H. Schmeisky, N. Hussain, S. Muhammad, M. Ibrahim and E. Safdar. 2008. Improvement of soil physical and chemical properties with compost application in rice-wheat cropping system *Pak. J. Bot.*, 40(1): 275-282.

Sarwar, G., N. Hussain, H. Schmeisky and S. Muhammad. 2007. Use of compost an environment friendly technology for enhancing rice-wheat production in Pakistan. *Pak. J. Bot.*, 39(5): 1553-1558.

Selvakumari, G., M. Baskar, D. Jayanthi and K.K. Mathan. 2000. Effect of integration of Flyash with fertilizers and organic manures on nutrient availability, yield and nutrient uptake of rice in alfisols. *J. Indian Soc. Soil Sci.*, 48: 268-278.

Sial, R.A., E.H. Chaudhary, S. Hussain and M. Naveed. 2007. Effect of organic manures and chemical fertilizers on grain yield of maize in rainfed area. *Soil & Environ.*, 26(2): 29-32.

Steel, R.G.D. and J.H. Torrie. 1980. Principles and Procedures of Statistics. 2nd Ed. McGraw Hill, Inc. Book Co. New York, USA.

Swarup, A. and N.P.S. Yaduvanshi. 2000. Effect of Integrated nutrient management on soil properties and yield of rice in Alkali soils. *J. Indian Soc. Soil Sci.*, 48: 279-282.

Terrance, D., M. Liebman, A. Cynthia, C.A. Cambardella and L. Richard. 2004. Maize response to composting and time of application of solid swine manure. *Agron. J.*, 96: 214-233.

Zahir, Z.A., M. Naveed, M.I. Zafar, H.S. Rehman, M. Arshad and M. Khalid. 2007. Evaluation of composted organic waste enriched with nitrogen and L-tryptophan for improving growth and yield of wheat (*Triticum aestivum* L.) *Pak. J. Bot.*, 39(5): 1739-1749.

(Received for publication 9 January 2008)