

ADDITION TO SEEDBORNE MYCOFLORA OF MUSTARD (*BRASSICA CAMPESTRIS* L.)

AMER-ZAREEN AND SHARFUNNAHAR

Department of Botany,
University of Karachi, Karachi-75270, Pakistan.

Mustard (*Brassica campestris* L.) is an important oil seed crop of Pakistan cultivated over 3,53,900 ha giving an yield of 2,85,600 tonnes annually (Anon., 1997). Richardson (1979, 1981, 1983) has listed 17 genera and 25 species of fungi from mustard seeds whereas a total of 9 genera and 17 species of fungi have been reported from mustard seeds in Pakistan (Iftikhar *et al.*, 1993). In the present study, 10 seed samples of mustard collected from different markets of Karachi were analyzed using Standard Blotter and Deep-Freezing methods (Anon., 1976). A total of 9 genera and 14 species of fungi viz., *Aspergillus flavus*, *A. niger*, *A. niveus*, *A. sulphureus*, *A. fumigatus*, *A. terreus*, *Chaetomium sp.*, *Cephaliophora irregularis*, *Curvularia lunata*, *Bipolaris hawaiiensis*, *Fusarium oxysporum*, *Penicillium sp.*, *Rhizopus stolonifer* and *Syncephalastrum sp.*, were isolated and identified (Table 1). Of these, *A. niger* and *A. fumigatus* have been reported from mustard seeds from India (Mondal *et al.*, 1988) and

Table 1. Incidence of fungi on mustard seeds.

Fungi	Standard Blotter Method			Deep Freezing Method				
	Infection %	Mean	SE ±	SD	Infection %	Mean	SE ±	SD
<i>Aspergillus flavus</i>	33.66	134.66	±37.07	90.8	12.34	160.75	±11.34	22.86
<i>A. niger</i>	1.51	6.16	±1.64	4.02	0.5	2	±1.35	2.7
<i>A. niveus</i>	1.87	7.5	±4.90	2.02	-	-	-	-
<i>A. sulphureus</i>	1.2	4.8	±2.05	5.03	2.81	11.25	±7.78	15.58
<i>A. fumigatus</i>	0.166	0.66	±0.32	0.8	-	-	-	-
<i>A. terreus</i>	-	-	-	-	0.68	2.75	±2.75	5.5
<i>Chaetomium sp.</i>	0.124	0.5	±0.49	1.22	0.25	1	±0.7	1.4
<i>Cephaliophora irregularis</i>	0.04	0.16	±0.166	0.408	-	-	-	-
<i>Curvularia lunata</i>	0.04	0.16	±0.166	0.408	-	-	-	-
<i>Bipolaris hawaiiensis</i>	-	-	-	-	0.25	1	±0.7	1.4
<i>Fusarium oxysporum</i>	-	-	-	-	0.75	3	±1.29	2.58
<i>Penicillium sp.</i>	0.083	0.33	±0.134	0.5	-	-	-	-
<i>Rhizopus stolonifer</i>	4.95	19.38	±8.49	20.8	7	28	±16.17	32.34
<i>Syncephalastrum sp.</i>	5.04	20.16	±11.90	29.15	5.87	23.5	±13	27.13

F. oxysporum from Pakistan (Richardson, 1979) whereas *A. niveus*, *A. sulphureus*, *A. terreus*, *Chaetomium* sp., *Cephaliophora irregularis*, *Curvularia lunata*, *Rhizopus stolonifer* and *Syncephalastrum* sp., do not appear to have been reported from mustard seeds (Richardson, 1979, 1981, 1983; Iftikhar *et al.*, 1993).

References

- Anonymous. 1976. International Rules for Seed testing. *Proc. Int. Seed Test. Assoc.*, 31:1-152
- Anonymous. 1997. *Agriculture Statistics of Pakistan*. Government of Pakistan. Ministry of Food, Agriculture & Livestock, Economic Wing, Islamabad, Pakistan.
- Iftikhar, A., I. Shamim and A. R. Bhatti. 1993. *Seed-borne microorganism in Pakistan. Check List 1991*. Pakistan Agriculture Research Council, Islamabad, Pakistan.
- Richardson, M. J. 1979. *An Annotated List of Seed-borne Diseases*. Int. Seed Test. Assoc., Zurich, Switzerland. 320 pp.
- Richardson, M. J. 1981. *An Annotated List of Seed-borne Diseases*. Supp. 1. Int. Seed Test. Assoc., Zurich, Switzerland. 78 pp.
- Richardson, M. J. 1979. *An Annotated List of Seed-borne Diseases*. Supp. 2. Int. Seed Test. Assoc., Zurich, Switzerland. 108 pp.
- Mondal, G. C., D. Nandi and B. Nandi. 1988. Effect of hydrolytic enzymes of storage fungi on seed deterioration. *Acta Agronomica Hungarica*, 36: 125-132.

(Received of publication 11 March 1999)