

## ENHANCEMENT IN GROWTH, NUTRIENT UPTAKE AND YIELD IN SALT STRESSED MAIZE BY FOLIAR APPLICATION OF METHIONINE

SADIA SHAHID<sup>1</sup>, ABIDA KAUSAR<sup>1\*</sup>, MUHAMMAD YASIN ASHRAF<sup>2</sup>,  
NOREEN AKHTAR<sup>1</sup> AND ZILL-I-HUMA NAZLI<sup>3</sup>

<sup>1</sup>Department of Botany, Government College Women University Faisalabad, 38000, Pakistan

<sup>2</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, Pakistan

<sup>3</sup>Department of Chemistry, Government College Women University Faisalabad, 38000, Pakistan

\*Corresponding author's email: [abidakausar@gcwuf.edu.pk](mailto:abidakausar@gcwuf.edu.pk)

### Abstract

Maize is leading cereal crop, having good quantity of vitamins, nutrients and minerals. It is cultivated all over the world for nourishment of human beings. Salinity is topmost problem of agricultural lands worldwide. Among numerous techniques being used in developing countries, use of organic compounds like amino acids is a cost effective technology. Amino acids have a valuable role in crop production and to reduce the changes produced due to abiotic stresses. In present experiment, four maize genotypes (FH1275, FH 936, FH 1231 and FH 1227) were used for two methionine levels (5 and 10 mg /L) under 80mM NaCl stress. After 120 days of propagating, four maize genotypes were evaluated for the measurements of various morphological and nutrient uptakes. Plants were uprooted and washed well. Yield parameters were examined at plant maturity, the methionine level 10 mg/ L showed better results as compared to 5 mg/ L both under saline and non-saline environments. Completely Randomized Design (CRD) with three factor factorial plan was used to assess the efficacy of amino acid on maize, under saline stress. Data was analyzed using suitable statistical analysis.

**Key words:** Nutrients uptake, Yield, Maize, Salinity, Methionine.

### Introduction

Maize in spite of being sensitive to salinity is nutritionally and economically very important (Estrada *et al.*, 2013). Maize (*Zea mays* L.) is a vital grain crop with 1053.8 million metric tons predictable global yield in 2019 (USDA, 2019). Due to high dietary values, global farming and low price, maize plays a key role for human and animal feedstuff (Tariq & Iqbal, 2010). Maize also contains a good quantity of sugar, proteins, fibers, fats as well as macro and micronutrients (Ali & Ashraf, 2011).

About one-third land area is affected by saline stress therefore adversely affecting the crop production in all over the world (United Nations, 2012). According to Panta *et al.*, (2014) salinity and sodicity ruins the agricultural land. Due to distortion of soil structure under salinity, plants cannot absorb nutrients accurately (Semida *et al.*, 2016). Saline stress causes a reduction in internode growth and leaf area resulting in obstruction of leaf initiation in maize (Sirhindi *et al.*, 2018). According to El Sayed (2011), initial growth of maize shoot is affected by salinity, whereas at later phases, growth is not affected harshly. It was also reported earlier that crop shoot and root growth as well as physiology are highly influenced under saline stress (Ashagre *et al.*, 2013).

Amino acids are very important as precursors of hormones, stress reducing mediators and nitrogen source (Maeda & Dudareva., 2012). In spite of the fact that amino acids are present in the soil naturally, but plants are not able to absorb them without the presence of any transporters in their roots (Khan *et al.*, 2020). Methionine, present in plant tissues and seeds, is an important sulfur containing essential amino acid (Galili & Amir, 2013) and its quantity limits the nutritional value of crop as it is comparable to base for proteinaceous food of animals and human beings (Galili *et al.*, 2016).

All the chemical processes occurring within the plants, are desperately affected by the existence of noxious ions in the soil (Ashraf & Tang, 2017) especially Na<sup>+</sup>/K<sup>+</sup> ratio due to boosted quantity of Na<sup>+</sup> ions under salinity, as plants absorbs more Na<sup>+</sup> and K<sup>+</sup> in salinity (Shahid *et al.*, 2011) which in turn results in enzymes inactivation (Sarwar & Shahbaz, 2020). Concentrations of ions in soil vary from each other under saline stress. For example, Ca<sup>2+</sup>, Na<sup>+</sup> and Cl<sup>-</sup> concentrations increase whereas Na<sup>+</sup> and K<sup>+</sup> concentrations exhibit a declining tendency in salinity (Akhtar *et al.*, 2003). It has been recorded in wheat that saline stress on physical characters is owed to augmented Na<sup>+</sup> and trivial K<sup>+</sup> concentration (Shamsi & Kobraee, 2013).

Osmotic stress and Na<sup>+</sup> toxicity in maize, grown under saline environment leads to meagre seed production (Schubert *et al.*, 2009). Kaya *et al.*, (2013) working on maize reported that total grain and ears production, weight of 1000 seed reduced under salinity.

### Materials and Methods

The experiment was performed using four genotypes of maize (FH 1275, FH 936, FH 1231 and FH 1227) to study the potential of salt tolerance using amino acid methionine (5 and 10 mg/L) under 80 mM NaCl stress.

The experiment was carried out at Government College Women University Faisalabad, Pakistan under natural wire house conditions. Ten seeds from each of the genotype were sown in plastic pots filled with sandy canal soil under 0 and 80 mM NaCl solution and three levels of methionine (5 and 10 mg/L) with three replicates of each treatment were maintained. After fifteen days of germination, the numbers of plants per pot was adjusted to four and were irrigated according to plant's requirement. Salinity was applied at four leaves stage and

maintained at 80 mM gradually. After seven days of salinity application, foliar spray of methionine with four days difference was applied. The pots were arranged in a Completely Randomized Design (CRD).

After 120 days of sowing, different growth and nutrients uptake parameters were recorded. Three plants from each treatment were removed at each harvest and washed with tap water. Water was blotted off from the surface of all plants for recording data. Shoot and root lengths (cm) with help of scale while shoot and root fresh weight (g), were measured at electrical balance.

Uptake of nutrients, e.g., sodium ( $\text{Na}^+$ ), potassium ( $\text{K}^+$ ), calcium ( $\text{Ca}^{2+}$ ) and chloride ( $\text{Cl}^-$ ), were calculated by using the method of Jackson (1962). The magnesium  $\text{Mg}^{2+}$  contents of shoot and root was determined according to the method in Handbook-60 by US Salinity Laboratory Staff, 1962. Yield parameters like number of kernels/ears, weight/ear (g), 100 seed weight (g) and 1000 seed weight (g) were measured. Statistically, data were analyzed by the three-way technique of analysis of variance and completely randomized design with three factors (maize, salinity, and methionine). The Least Significant Difference (LSD) test at 5% probability was used to calculate the differences among the mean values.

## Results

### Growth parameters

**Shoot and root lengths:** Genotype FH 1275 of maize maintained the highest shoot length (60.3 cm) and least value was documented in FH 1227 (38.4 cm) in control. Peak value of plant height was noted under methionine 10 mg/L (72.3 cm), while minimum under non saline conditions (53.5 cm). The overall decrease in shoot lengths was 39.9% under salinity in FH 1275 in saline stress as compared to the normal environment. (Fig. 1, Table 1). Salinity has highly influenced the plant root length (23.2 cm) as compared to non-saline condition (36.5 cm) showing an overall decrease in root length, 36.4% in FH1275. Application of 10 mg/L methionine showed maximum (32 cm) value in FH1275 while lowest value of root length was noted in FH 1227 (21.5 cm) under 80 mM NaCl. Overall increase in root length under 10mg/L methionine and 80Mm NaCl was 39.13% in FH 1275 (Fig. 1, Table 1).

**Shoot and root fresh and dry weights:** Highest shoot fresh weight (81.8 g) was observed in FH 1275 whereas minimum (50.4g) in FH 1227 in control plants. Whereas the value of 60.4 (g) in FH 1275 and 21.4 (g) in FH 1227 was measured in saline environments. Overall decrease in shoot fresh weight was 26.1 % under salinity in FH 1275. Spray of methionine 10 mg/L (105.0 g) on FH 1275, showed the highest whereas the lowest shoot fresh weight (64.3 g) reading was observed in FH 1227 under control conditions. Overall increase (22.5%) in shoot fresh weight under 10mg/L methionine in all maize genotypes was recorded. (Fig. 1, Table 1). Maximum root fresh weights was shown by FH 1275 (9.45 g), whereas FH 1227 performed poorly with minimum (4.8 g) root fresh weight in control medium. However, in saline medium the values

7.89 (g) in FH 1257 and lowest in FH1227 were 1.8 (g) were investigated. Overall decrease in root fresh weight was 17.4 % under salinity in all maize genotypes were recorded. Application of methionine 10 mg/L was depicted as most effective dose concentration with highest value of root fresh weight (13.8 g) in FH 1275 and lowest (6.95 g) was observed in FH 1227. Percent increase in root fresh weight in FH 1275 was 34.6 % under 10mg/L methionine and 80mM saline stress (Fig. 1, Table 1).

### Analysis of Nutrients

**Shoot and root  $\text{Na}^+$ :** Amount of  $\text{Na}^+$  was found elevated in shoot under salt stress (16.8 mg/g D. W.) (D.W = Dry weight) than control (6.8 mg/g D. W.) conditions (Fig. 1). Genotype FH 1227 maintained the highest (27.39 mg/g D. W.) while the FH 1275 showed minimum value of  $\text{Na}^+$  (5.3 mg/g D. W.) in shoots suggesting overall rise in shoot  $\text{Na}^+$  36.6% in FH 1227 under salinity. Highest value of shoot  $\text{Na}^+$  (7.1 mg/g D. W.) in FH 1227 (28.30% decrease) whereas lowest value of shoot  $\text{Na}^+$  (3.8 mg/g D. W.) was recoded in FH 1275 (26.3% decrease) under 10mg/L methionine (Fig. 1). Highest value of root  $\text{Na}^+$  (27.3 mg/g D. W.) (103.7% increase) was documented in FH 1227 under salinity while lowest value (16.8 mg/g D. W.) (147% increase) was measured in FH 1227 under saline stress. Highest value of root  $\text{Na}^+$  (16.77 mg/g D. W.) (62.7% increase) in FH 1227 while lowest value (9.7 mg/g D. W.) (73.19 % increase) in FH 1275 was observed by the foliar spray of 10mg/L methionine (Fig. 2, Table 1).

**Shoot and root  $\text{K}^+$ :** Saline stress minimized root  $\text{K}^+$  to 17.7 mg/g D. W. compared to control (21.3 mg/g D. W.) in FH 1275 (16.9% decrease) while FH 1227 showed lowest value 7.7 mg/g D. under saline stress as compared to control (15.8 mg/g D. W.) (51.2% decrease). Genotype FH 1275 was observed to achieve high value (20.8 mg/g D. W.) (17.5% increase) of shoot  $\text{K}^+$  by the spray of methionine 10 mg/L whereas the lowest value (11.5 mg/g D. W.) (49.3% increase) was noted by FH 1227 under saline stress (Fig. 3, Table 1). FH 1275 maintained the highest value of root  $\text{K}^+$  (7.8 mg/g D. W.) (34.4% decrease) under saline stress of 80 mM whereas FH 1227 showed least value (5.06 mg/g D. W.) (41.8% decrease) as compared to control. Foliar spray of methionine 10 mg/L resulted in highest root  $\text{K}^+$  (12.2 mg/g D. W.) (54.1% increase) by FH 1275 while lowest value was noted in FH 1227 (7.9 mg/g D. W.) (58% increase) under salt stress (Fig. 3, Table 1).

**Shoot and root  $\text{Ca}^{2+}$ :** Substantial drop (3.8 mg/g) (44.8% decrease) of shoot  $\text{Ca}^{2+}$  was noted in genotype FH 1227 however maximum value (6.29 mg/g) (37.4% decrease) of  $\text{Ca}^{2+}$  was documented in FH 1275 under saline stress of 80mM NaCl compared to control (Fig. 2, Table 2). Maximum (11.4 mg/g) (83.8% increase) plant root  $\text{Ca}^{2+}$  content was observed in FH 1275 and lowest value (6.39 mg/g) (65.7% increase) was noted in FH 1227 under saline condition by methionine 10 mg/L (Fig. 2, Table 2).




Fig. 1. Effect of methionine (5 and 10 mg/L) on shoot and root lengths (cm) and shoot and root fresh weights (g) and shoot Na<sup>+</sup> (mg/g D. W.) of four maize genotypes (FH 1275, FH 936, FH 1231 and FH 1227) under 80 mM NaCl stress.




Fig. 2. Effect of methionine (5 and 10 mg/L) on root Na<sup>+</sup> (mg/g dry W.), shoot and root Ca<sup>2+</sup> (mg/g D. W.) and shoot Cl<sup>-</sup> (mg/g D. W.) of four maize genotypes (FH 1275, FH 936, FH 1231 and FH 1227) under 80 mM NaCl stress.

**Shoot and root Cl<sup>-</sup>:** Salinity increased amount of Cl<sup>-</sup> (16.5 mg/g) in shoots of FH 1275 compared to control condition (6.6 mg/g) (150% increase) while value of shoot Cl<sup>-</sup> (17.8 mg/g) was observed in FH 1227 under salinity of 80mM as compared to control (8.1 mg/g) (120% increase). FH 1227 maintained the highest value of shoot Cl<sup>-</sup> (17.03 mg/g) (4.3% decrease) while lowest value was observed in FH 1275 (15.23 mg/g) (7.8% decrease) under 5mg/L

methionine spray and 80Mm salinity (Fig. 2, Table 2). FH 1227 maintained the peak value (21.6 mg/g) (87.8% increase) of root Cl<sup>-</sup> while lowest value was observed by FH 936 (12.1 mg/g) (108.6% increase) under saline stress of 80Mm. FH 1227 showed maximum value of root Cl<sup>-</sup> (19.2 mg/g) (24.5% decrease) and FH 936 showed lowest value root Cl<sup>-</sup> (7.4 mg/g) (41.3% decrease) under 10mg/L spray of methionine. (Fig. 3, Table 2).

**Table 1. Mean squares values from analysis of variance of data for shoot and root lengths, fresh weights and shoot and root Na<sup>+</sup> and K<sup>+</sup> as affected by foliar application of methionine (5 and 10 mg/L) under 80 mM (NaCl) salt stress.**

Source	Mean squares								
	df	Shoot length	Root length	Shoot fresh weight	Root fresh weight	Shoot Na <sup>+</sup>	Root Na <sup>+</sup>	Shoot K <sup>+</sup>	Root K <sup>+</sup>
Varieties (V)	3	1086.96***	570.80***	6474.5***	143.881***	47.6565***	231.716***	277.114***	66.850 ***
Salinity (S)	1	3074.59***	2134.00***	13750.9***	167.964***	72.4240***	872.506***	595.700***	298.901***
Treatment (T)	2	670.79**	502.00***	1373.6***	51.731***	28.5384***	191.797***	182.657***	98.845**
V × S	3	9.85 <sup>ns</sup>	1.16 <sup>ns</sup>	35.1 <sup>ns</sup>	0.248 <sup>ns</sup>	1.2503 <sup>ns</sup>	11.493 <sup>ns</sup>	1.775 <sup>ns</sup>	1.526 <sup>ns</sup>
V × T	6	26.37 <sup>ns</sup>	12.82 <sup>ns</sup>	2.4 <sup>ns</sup>	1.274 <sup>ns</sup>	0.6203 <sup>ns</sup>	4.600 <sup>ns</sup>	1.354 <sup>ns</sup>	1.891 <sup>ns</sup>
S × T	2	83.90**	22.38 <sup>ns</sup>	60.6*	0.938 <sup>ns</sup>	0.3327 <sup>ns</sup>	114.990***	40.577**	5.877 <sup>ns</sup>
V × S × T	6	12.12 <sup>ns</sup>	2.73 <sup>ns</sup>	46.3*	0.474 <sup>ns</sup>	0.2838 <sup>ns</sup>	3.371 <sup>ns</sup>	3.700 <sup>ns</sup>	1.372 <sup>ns</sup>
Error	48	16.12	13.61	20.0	3.701	2.6472	6.009	7.016	5.194
<b>Total</b>	<b>71</b>								

\*\*\*=  $p < 0.001$ ; \*\*=  $p < 0.01$ ; \*=  $p < 0.05$ ; ns= Non-significant**Table 2. Mean squares values from analysis of variance of data for shoot and root Ca<sup>2+</sup>, Cl<sup>-</sup>, Mg<sup>2+</sup> as affected by foliar application of methionine (5 and 10 mg/L) under 80 mM (NaCl) salt stress.**

Source	Mean squares						
	Df	Shoot Ca <sup>2+</sup> (mg/g D. W.)	Root Ca <sup>2+</sup> (mg/g D. W.)	Shoot Cl <sup>-</sup> (mg/g D.W.)	Root Cl <sup>-</sup> (mg/g D.W.)	Shoot Mg <sup>2+</sup> (mg/g D.W)	Root Mg <sup>2+</sup> (mg/g D. W.)
Genotypes (G)	3	63.902***	134.800***	207.919***	283.110***	34.6474***	25.8914***
Salinity (S)	1	19.709*	21.725*	3.754 <sup>ns</sup>	5.917 <sup>ns</sup>	23.8856**	8.1541***
Treatment (T)	2	127.011***	104.978***	383.763***	503.233***	67.0926***	41.9845***
G × S	3	3.085 <sup>ns</sup>	0.042 <sup>ns</sup>	1.479 <sup>ns</sup>	0.211 <sup>ns</sup>	0.3946 <sup>ns</sup>	1.0926 <sup>ns</sup>
G × T	6	0.341 <sup>ns</sup>	1.417 <sup>ns</sup>	14.661*	18.907**	0.0427 <sup>ns</sup>	0.2147 <sup>ns</sup>
S × T	2	27.237**	28.257**	45.260***	76.230***	22.3094**	14.5723***
G × S × T	6	0.080 <sup>ns</sup>	2.378 <sup>ns</sup>	0.860 <sup>ns</sup>	1.015 <sup>ns</sup>	0.1006 <sup>ns</sup>	0.5209 <sup>ns</sup>
Error	48	3.840	3.567	5.302	4.508	3.2138	0.5235
<b>Total</b>	<b>71</b>						

\*\*\*=  $p < 0.001$ ; \*\*=  $p < 0.01$ ; \*=  $p < 0.05$ ; ns= Non-significant**Table 3. Analysis of variance (mean squares) of data for weight of ear/plant, number of ears /plants, number of kernels/ cobs, 100 kernels weight (g) and 1000 kernels weight (g) as affected by foliar application of methionine (5 and 10 mg/L) under 80 mM (NaCl) salt stress.**

Source	Mean squares					
	Df	Weight of ear/ plant	Number of ears/ plants	Number of kernels/ cobs	100 kernels weight (g)	1000 kernels weight (g)
Varieties (V)	3	356540***	33303***	239633***	549.743***	495.843***
Salinity (S)	1	6506***	27643***	149058***	13.390*	14.405**
Treatment (T)	2	261 ns	540074**	38314***	7.229*	3.112*
V × S	3	6706***	63221**	23816***	33.793***	30.453**
V × T	6	268 ns	4876321ns	6007***	2.907 <sup>ns</sup>	1.0654ns
S × T	2	4 ns	33564ns	25985***	39.154***	43.415**
V × S × T	6	4 ns	3465**	2675***	25.209***	19.3354***
Error	48	243	274	165	1.950	1.8864
<b>Total</b>	<b>71</b>					

\*\*\*=  $p < 0.001$ ; \*\*=  $p < 0.01$ ; \*=  $p < 0.05$ ; ns= Non-significant

**Shoot and root Mg<sup>2+</sup>:** The maximum plant shoot Mg<sup>2+</sup> content (5.44 mg/g D. W.) was observed in FH 1275 (28% decrease) while the lowest value (2.4 mg/g D. W.) (47.8% decrease) was noted in FH 1227 under saline conditions compared to control. Foliar application 10mg/L methionine resulted in highest value of shoot Mg<sup>2+</sup> in FH 1275 (8.17 mg/g D. W.) (24.6% increase) while lowest value was observed in FH1227 (4.3 mg/g D. W.) (38.7% increase) (Fig. 3, Table 2). Root Mg<sup>2+</sup> under saline conditions showed maximum (4.8 mg/g D. W.) (27 % decrease) value and lowest value (2.4 mg/g D. W.) (44.18 % decrease) was observed in FH 1227 as compared to control. Genotype FH 1275 was recoded to accomplish high value (6.7 mg/g D. W.) (24.0 % increase) of

plant root Mg<sup>2+</sup> with spray of methionine 10 mg/L whereas the lowest value (3.6 mg/g D. W.) (24.1% increase) was noted in FH 1227 (Fig. 3, Table 2).

**Yield parameters:** Maximum value of weight of ear (180 g) (51.3% decrease) in FH1275 and minimum was observed in FH-1227 (56.6 g) (41.6% decrease) under the effect of salinity as compared to control. The maximum weight by FH-1275 (256 g) (42.2% increase) under the treatment of methionine 10 mg/L was recorded whereas the lowest value was calculated by FH-1227 (190 g) (235% increase) compared to saline stress of 80 mM (Fig. 4, Table 3).




Fig. 3. Effect of methionine (5 and 10 mg/L) on root Cl<sup>-</sup> (mg/g D.W.), shoot and root K<sup>+</sup> (mg/g D.W.) and shoot and root Mg<sup>2+</sup> (mg/g D. W.) of four maize genotypes (FH 1275, FH 936, FH 1231 and FH 1227) under 80 mM NaCl stress.




Fig. 4. Effect of methionine (5 and 10 mg/L) on weight of ear/plant (g), number of kernels /cob, number of ears/plant, 100 kernels weight (g) and 1000 kernels weight (g) (mg/g D.W.) of four maize genotypes (FH 1275, FH 936, FH 1231 and FH 1227) under 80 mM NaCl stress.

Maximum value of number of ears/plant (2.03) (44.5% decrease) in FH 1275 and minimum was observed in FH-1227 (66) under the effect of salinity as compared to control (.01% decrease). The maximum number of ears/plants was recorded by FH-1275 (2.3g) (13.7% increase) under the treatment of methionine 10 mg/L whereas the lowest value was calculated by FH-1227 (1.3) (100 % increase) as compared to saline stress of 80 mM (Fig. 4, Table 3).

Peak value of number of kernels/cob (480) (5.6 % decrease) in FH 1275 and minimum was observed in FH-1227 (156) (20.4 % decrease) under the effect of salinity compared to control. The maximum number of kernels/cobs was recorded by FH-1275 (546) (2.82% increase) under the treatment of methionine 10 mg/L whereas the lowest value was calculated by FH-1227 (390) (100% increase) as compared to saline stress of 80 mM (Fig. 4, Table 3).

Highest value of 100 kernels weight (12 g) (50% decrease) in FH 1275 and minimum was observed in FH-1227 (0.9) (15.9% decrease) under the effect of salinity compared to control. The maximum weight was recorded by FH-1275 (10g) (16.6% decrease) under the treatment of methionine 10 mg/L whereas the lowest value was calculated by FH-1227 (0.59 g) (34.4 % increase) as compared to saline stress of 80 mM (Fig. 4, Table 3).

Peak value of 1000 kernels weight (95 g) (51% decrease) in FH 1275 and minimum was observed in FH-1227 (50) (47.3% decrease) under the effect of salinity compared to control. The maximum weight was recorded by FH-1275 (210g) (121% increase) under the treatment of methionine 10 mg/L whereas the lowest value was calculated by FH-1227 (30) (40 % increase) as compared to saline stress of 80 mM (Fig. 4, Table 3).

## Discussion

Saline soils adversely affect plant growth and development throughout in the world. Various studies demonstrated the negative impacts of salinity on several plant morphological and biochemical traits (Hussein *et al.*, 2017; Di Mola *et al.*, 2017; Marium *et al.*, 2021). Salinity reduced the leaf surface expansion and biomass of purslane (*Portulaca oleracea* L.) (Alam *et al.*, 2015). The relative shoot length of seedlings of salt tolerant sorghum plants, enhanced the biomass production of plants and increased the absorption of  $K^+$  while lowered the shoot  $Na^+$  contents (Krishnamurthy *et al.*, 2007). Exogenous application of methionine (5 and 10 mg/L) enhanced the salinity tolerance of maize genotypes in their morphological parameters, nutrients uptake and yield features. Present results depicted that foliar application of methionine significantly enhanced the root and shoot lengths, fresh and dry weights of all four maize genotypes. Similar results were presented by Hammad & Ali, (2014) and Zaki *et al.*, (2007) showing that bio stimulants i.e., amino acids when applied then amplified all developmental traits of wheat. Earlier investigations showed that salinity badly effects plant biomass and productivity which results in the less crop yield because

number of soluble salts in soil and water, increased (Ahanger *et al.*, 2014; Flowers *et al.*, 2015). Khan *et al.*, (2014) reported improvement in plant growth upon the spray of methionine with and without saline environments, suggesting ethylene formation under salt stress. The increased methionine level leads to the production of osmolytes as glycine betaine or other antioxidants and reduces ethylene production in salinity as depicted in current studies and found in mung bean (*Vigna radiata* L (El-Zohiri & Asfour, 2009). It was proved that methionine enhanced the tolerance potential of plants maize and Faba beans (Sh Sadak *et al.*, (2015).

In present study the absorbance of nutrients is badly affected by salinity, but the methionine meaningfully enhanced the nutrients uptake ( $K^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ ). It may be due to presence of antagonistic role of toxic ions. According to Wang *et al.*, (2012), the plants that are grown under salinity, the absorbance of essential nutrients ( $K^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$ ) is highly affected by competition with the other lethal ions like  $Na^+$  and  $Cl^-$  (Kausar & Gull, 2019). All maize genotypes, in present experiment, gathered a significant extent of shoot plus root  $Na^+$  and  $Cl^-$  in saline environment but genotype FH 1231 and FH 1227 enhanced the amount of these toxic nutrients as compared to FH 1275 and FH 936. Wang *et al.*, (2012) described that enhanced level of root and shoot  $Na^+$  and  $Cl^-$  have been ascribed to the struggle with other nutrients like  $K^+$ ,  $Ca^{2+}$  and  $Mg^{2+}$ . Similar observations were noticed in tomato by Tantawy *et al.*, (2009) and in faba beans by Sh Sadak *et al.*, (2015) that foliar spray of amino acid resulted in increased amount of  $Ca^{2+}$ ,  $P^{2+}$ ,  $K^+$  and  $K^+/Na^+$  in faba bean. In present experiment the enhanced amounts of  $K^+$ ,  $N^+$ ,  $Mg^{2+}$  and  $Ca^{2+}$  stored in shoot and root resulted in better yield of the crop which were similar with the study of Nahed *et al.*, (2010) who worked on *Thuja orientalis* and suggested that N, P, K (%) was increased by amino acid spray on the plant.

Salinity drastically reduced the crop yield in several crops (Ashraf, 2010) like maize (Ali *et al.*, 2008), wheat (Ranjbar, 2010), sorghum (kausar & Gull, 2019) and in cucumber (Marium *et al.*, 2021). Similar outcomes were observed in present trial where NaCl decreased the number of ears on each maize plant and regular yield, but the foliar application of methionine reduced the fatal influence of salinity on all maize genotypes. However, genotypes FH 1275 and FH 936 exhibited more tolerance to salinity by better maintenance of different yield attributes than the other genotypes (FH 1231 and FH 1227). According to Muneer *et al.*, (2014) while working on *Solanum lycopersicum* noticed in the reduction of plant yield due to low rate of photosynthesis under saline environment. According to Shahbaz *et al.*, (2012) working on different vegetables showed an increase in crop yield under foliar spray of osmo-protectants. It has been depicted from the results that FH 1275 and FH 936 exhibited better performance under saline and non-saline conditions by the foliar application of 10mg/L of methionine compared to FH 1231 and FH 1227. Moreover 10mg/L foliar spray enhanced the plant growth and nutrients uptake which may be due to activation of salt tolerant genes through the signaling effect of methionine.

## Conclusion

Methionine enhanced the plant biomass and nutrient uptake of all maize genotypes both under salinity and normal conditions. It is concluded that FH 1275 and FH 936 and 10 mg/L level of methionine foliar spray showed better results as compared to the FH 1231 and FH 1231 and could be recommended to farmers to grow in saline soils to increase growth and yield of maize genotypes in salt hit areas of the world.

## References

- Ahanger, M.A., A. Hashem, E.F. Abd-Allah and P. Ahmad. 2014. Arbuscular mycorrhiza in crop improvement under environmental stress. In Emerging technologies and management of crop stress tolerance. *Acad. Press.*, 69-95.
- Akhtar, S., A. Wahid and E. Rasul. 2003. Emergence, growth and nutrient composition of sugarcane sprouts under NaCl salinity. *Biol. Plant.*, 46: 113-116.
- Alam, M., A.S. Juraimi, M.Y. Rafii and A. Abdul Hamid. 2015. Effect of salinity on biomass yield and physiological and stem-root anatomical characteristics of purslane (*Portulaca oleracea* L.) accessions. *Biomed. Res. Int.*, 2015.
- Ali, Q. and M. Ashraf. 2011. Induction of drought tolerance in maize (*Zea mays* L.) due to exogenous application of trehalose: growth, photosynthesis, water relations and oxidative defense mechanism. *J. Agron. Crop Sci.*, 197: 258-271.
- Ali, Q., M. Ashraf, M. Shahbaz and H. Humera. 2008. Ameliorating effect of foliar applied proline on nutrient uptake in water stressed maize (*Zea mays* L.) plants. *Pak. J. Bot.*, 40: 211-219.
- Ashagre, H., M. Shelema, R. Kedir and S. Ebsa. 2013. Seed germination and seedling growth of haricot bean (*Phaseolus vulgaris* L.) cultivars as influenced by copper sulphate. *World J. Agric. Sci.*, 1: 312-317.
- Ashraf, M. 2010. Registration of 'S-24' Spring Wheat with Improved Salt Tolerance. *J. Plant Regist.*, 4: 34-37.
- Ashraf, U. and X. Tang. 2017. Yield and quality responses, plant metabolism and metal distribution pattern in aromatic rice under lead (Pb) toxicity. *Chemosphere.*, 176: 141-155.
- Di Mola, I., Y. Roupheal, G. Colla., M. Fagnano., R. Paradiso and M. Mori. 2017. Morphophysiological traits and nitrate content of greenhouse lettuce as affected by irrigation with saline water. *Hort. Sci.*, 52: 1716-1721.
- El Sayed, H.E.S.A. 2011. Influence of salinity stress on growth parameters, photosynthetic activity and cytological studies of *Zea mays* L. plant using hydrogel polymer. *Agric. Biol. J. N. Ame.*, 2: 907-920.
- El-Zohiri, S.S.M. and Y.M. Asfour. 2009. Effect of some organic compounds on growth and productivity of some potato cultivars. *Ann. Agric. Sci. Moshtohor.*, 47: 403-415.
- Estrada, B., R. Aroca, F.J. Maathuis., J.M. Barea and J.M. Ruiz-lozano. 2013. Arbuscular mycorrhizal fungi native from a Mediterranean saline area enhance maize tolerance to salinity through improved ion homeostasis. *Plant Cell Environ.*, 36: 1771-1782.
- Flowers, T.J., R. Munns and T.D. Colmer. 2015. Sodium chloride toxicity and the cellular basis of salt tolerance in halophytes. *Ann. Bot.* 115: 419- 431.
- Galili, G. and R. Amir. 2013. Fortifying plants with the essential amino acids lysine and methionine to improve nutritional quality. *Plant Biotechnol. J.*, 11: 211-222.
- Galili, G., R. Amir and A.R. Fernie. 2016. The regulation of essential amino acid synthesis and accumulation in plants. *Ann. Rev. Plant Biol.*, 67: 153-178.
- Hammad, S.A. and O.A. Ali. 2014. Physiological and biochemical studies on drought tolerance of wheat plants by application of amino acids and yeast extract. *Ann. Agric. Sci.*, 59: 133-145.
- Hussein, M., A. Embiale, A. Husen., I.M. Aref and M. Iqbal. 2017. Salinity-induced modulation of plant growth and photosynthetic parameters in faba bean (*Vicia faba*) cultivars. *Pak. J. Bot.*, 49: 867-877.
- Jackson, M.L. 1962. Interlayering of expansible layer silicates in soils by chemical weathering. *Clays & Clay Miner.*, 11: 29-46.
- Kausar, A. and M. Gull. 2019. Influence of salinity stress on the uptake of magnesium, phosphorus, and yield of salt susceptible and tolerant sorghum cultivars (*Sorghum bicolor* L.). *J.A.B.B.*, 7: 53-58.
- Kaya, C., M. Ashraf, M. Dikilitas and A.L. Tuna. 2013. Alleviation of salt stress-induced adverse effects on maize plants by exogenous application of indoleacetic acid (IAA) and inorganic nutrients-A field trial. *Aust. J. Crop Sci.*, 7: 249.
- Khan, N.A., M.I.R. Khan, M. Asgher., M. Fatma., A. Masood and S. Syeed. 2014. Salinity tolerance in plants: revisiting the role of sulfur metabolites. *Plant Physiol. Biochem.*, 2: 2.
- Khan, N., S. Ali, P. Zandi, A. Mehmood, S. Ullah, M. Ikram, M.A.S. Ismail and M. Babar. 2020. Role of sugars, amino acids and organic acids in improving plant abiotic stress tolerance. *Pak. J. Bot.*, 52: 355-363.
- Krishnamurthy, L., R. Serraj, C.T. Hash, A.J. Dakheel and B.V. Reddy. 2007. Screening sorghum genotypes for salinity tolerant biomass production. *Euphytica.*, 156: 15-24.
- Maeda, H. and N. Dudareva. 2012. The shikimate pathway and aromatic amino acid biosynthesis in plants. *Ann. Rev. Plant Biol.*, 63: 73-105.
- Marium, A., A. Kausar, M.Y. Ashraf, H. Kanwal and Z.I.H. Nazli. 2021. Enhancement in biomass and nutrients uptake in cucumber through methionine and phenylalanine under saline soils. *Pak. J. Agric. Sci.*, 58: 1.
- Muneer, S., Y. Park, A. Manivannan, P. Soundararajan and B. Jeong. 2014. Physiological and proteomic analysis in chloroplasts of *Solanum lycopersicum* L. under silicon efficiency and salinity stress. *Int. J. Mol. Sci.*, 15: 21803-21824.
- Nahed, G., A.A. Abdel Aziz., M. Mazher and M.M. Farahat. 2010. Response of vegetative growth and chemical constituents of *Thuja orientalis* L. plant to foliar application of different amino acids at Nubaria. *Am. J. Sci.*, 6: 295-301.
- Panta S., T. Flowers, P. Lane., R. Doyle and G. Haros. 2014. Halophytes agriculture: success stories. *Environ. Exp. Bot.*, 107: 71-83.
- Ranjhar, G.H. 2010. Salt sensitivity of two wheat cultivars at different growth stages. *World Appl. Sci. J.*, 11: 309-314.
- Sarwar, Y. and M. Shahbaz. 2020. Modulation in growth, photosynthetic pigments, gas exchange attributes and inorganic ions in sunflower (*Helianthus annuus* L.) by strigolactones (GR24) achene priming under saline conditions. *Pak. J. Bot.*, 52: 23-31.
- Schubert, S., A. Neubert., A. Schierholt., A. Sümer and C. Zörb. 2009. Development of salt-resistant maize hybrids: the combination of physiological strategies using conventional breeding methods. *Plant Sci. J.*, 177: 196-202.
- Semida, W.M., T.A. Abd El-Mageed., S.A. Howladar and M.M. Rady. 2016. Foliar-applied alpha-tocopherol enhances salt-tolerance in onion plants by improving antioxidant defense system. *Aust. J. Crop Sci.*, 10: 1030.
- Sh Sadak, M., M.T. Abdelhamid and U. Schmidhalter. 2015. Effect of foliar application of aminoacids on plant yield and some physiological parameters in bean plants irrigated with seawater. *Acta Biol. Colomb.*, 1: 141-152.

- Shahbaz, M., M. Ashraf, F. Al-Qurainy and P.J. Harris. 2012. Salt tolerance in selected vegetable crops. *Crit. Rev. Plant Sci.*, 31: 303-320.
- Shahid, M.A., M.A. Pervez, R.M. Balal, N.S. Mattson, A. Rashid, R. Ahmad, C.M. Ayyub and T. Abbas. 2011. Brassinosteroid (24-epibrassinolide) enhances growth and alleviates the deleterious effects induced by salt stress in pea (*Pisum sativum* L.). *Aust. J. Crop Sci.*, 5: 500.
- Shamsi, K. and S. Kobraee. 2013. Biochemical and physiological responses of three wheat cultivars (*Triticum aestivum* L.) to salinity stress. *Ann. Biol. Res.*, 4: 180-185.
- Sirhindi, G., R. Bhardwaj, M. Kumar, S. Kumar, N. Dogra, H. Sekhon, S. Kaushik and I. Madaan. 2018. Physiological roles of brassinosteroids in conferring temperature and salt stress tolerance in plants. In *Metabolic Adaptations in Plants During Abiotic Stress* CRC Press. 339-358.
- Tantawy, A.S., A.M.R. Abdel-Mawgoud, M.A. El-Nemr and Y.G. Chamoun. 2009. Alleviation of salinity effects on tomato plants by application of amino acids and growth regulators. *Eur. J. Sci. Res.*, 30: 484-494.
- Tariq, M. and H. Iqbal. 2010. Maize in Pakistan-an overview. *Kasetsart J. Nat. Sci.*, 757-763.
- USDA, F., 2019. EU Biofuels Annual. *GAIN Report* NL9022.
- Wang, Y., L. Li, W. Cui., W., S. Xu., W. Shen and R. Wang. 2012. Hydrogen sulfide enhances alfalfa (*Medicago sativa*) tolerance against salinity during seed germination by nitric oxide pathway. *Plant Soil.*, 351: 107-119.
- Zaki, M.S., M. Nabila, Hassanein and K.M. Gamal El-Din. 2007. Growth and yield of some wheat cultivars irrigated with saline water in newly cultivated land as affected by biofertilization, *Res. J. Appl. Sci.*, 3: 1121-1126.

(Received for publication 22 January 2020)