

## EFFICACY AND PERSISTENCE OF MICROBIAL ANTAGONISTS AGAINST *SCLEROTIUM ROLFSII* UNDER FIELD CONDITIONS

FOUZIA YAQUB\* AND SALEEM SHAHZAD

Pest & Disease Research Lab., Department of Agriculture and Agribusiness Management,  
University of Karachi, Karachi-75270, Pakistan

\*E-mail: fouziayaqub\_biologist@yahoo.com

### Abstract

*Sclerotium rolfii* showed significant negative effect on plant growth due to severe root colonization, whereas, presence of the microbial antagonists showed significant positive effect on plant growth by reducing the colonization of roots by *S. rolfii*. Highest root colonization by *S. rolfii* and significant reduction in plant growth were observed in sunflower and mungbean plants growing in soil artificially infested with sclerotia of *S. rolfii*. Use of biocontrol agents in *S. rolfii* infested soil showed significant reduction in Root Colonization Index accompanied by increase in plant growth. *Bradyrhizobium* sp., was found most effective ( $p < 0.01$ ) followed by *Rhizobium* sp., *Trichoderma harzianum*, *T. pseudokoningii*, *T. polysporum* and *T. virens*.

Efficacy of biocontrol agents was comparatively suppressed in *S. rolfii* infested soil as compared to non-infested soil. There was no significant difference in plant length in infested and non-infested soils where biocontrol agents were used. However, plant weight was less in infested soils as compared to non-infested soils. The microbial antagonists persisted in the soil and produced similar results when mungbean and sunflowers were re-sown in the same plots.

### Introduction

*Sclerotium rolfii* Sacc., [teleomorph: *Athelia rolfii* (Curzi) Tu & Kimbrough] is an economically important soil-borne pathogen that is very common in tropical, sub-tropical and other warm temperate regions of the world. The pathogen causes root rot, stem rot, wilt and foot rot diseases on more than 500 species of cultivated and wild plants including almost all the agricultural and horticultural crops (Aycock, 1966; Punja, 1985; Domsch *et al.*, 1980; Farr *et al.*, 1989; Cilliers *et al.*, 2000; Ciancio & Mukerji, 2007). Mostly *S. rolfii* diseases have been reported on dicotyledonous hosts, but several monocotyledonous species have also been infected (Aycock, 1966; Ciancio & Mukerji, 2007). Humid weather is conducive to sclerotial germination and mycelial growth. Consequently the diseases caused by the fungus are more serious in tropical and subtropical regions than in temperate regions (Yorinori, 1994). The large number of sclerotia produced by *S. rolfii* and their ability to persist in the soil for several years, as well as the profuse growth rate of the fungus make it well suited facultative parasite and a pathogen of major importance throughout the world (Punja, 1988).

The first confirmed report of losses due to the pathogen in USA was made by Rolfs (1892) on tomato (*Lycopersicon esculentum* Miller) in Florida. Aycock (1966) estimated that *S. rolfii* produced yield depletion ranging from 1-60% in different peanut fields in southern peanut growing region of USA resulting in losses of up to US\$ 10-20 million. The first report of *S. rolfii* from Pakistan was made by Ahmed *et al.*, (1984) who isolated it from maize (*Zea mays* L.). Subsequent reports were made from oat (*Avena sativa* L.) and mash bean (*Vigna mungo* L.) Hepper by Shahzad & Ghaffar (1995), lentil (*Lens culinaris* L.) Medic. by Iqbal *et al.*, (1995), apple (*Malus sylvestris* L.) by Jahangir *et al.*, (1995), and seeds of sugarbeet (*Beta vulgaris* L.) by Ruqia (2001).

Use of chemical fungicides is no doubt the most effective method for the control of plant diseases. However, in view of the complexities arising from the use of chemical pesticides, such as harmful effects on

environment and non-target organisms including man, domestic animals, beneficial insects and wild life, efforts have been focused to develop alternative approaches which are safe for all stakeholders i.e. humans, animals, environments and crops (Atlas & Bartha, 1998). Of the various methods used, the use of microorganism as biocontrol agents has provided a very promising alternative and less hazardous method for plant disease control (Papavizas & Lumsden, 1980).

Diseases caused by *S. rolfii* continue to receive considerable attention with regard to the development of biological control strategies (Tjamos *et al.*, 1992). The application of fungi as biological control agents, especially *Trichoderma* spp., to control *S. rolfii* has been attempted (Henis, 1984; Papavizas & Lewis, 1989). *T. harzianum* reduced root rot of sugar beets (Ciccarese *et al.*, 1992), stem rot of ground nut (Cilliers *et al.*, 2000), damping-off and stem rot of cowpea (Adandonon, 2000; Kossou *et al.*, 2001), root rot of beans and tomatoes (Elad *et al.*, 1980), collar rot and seedling death of lentil (Agrawal *et al.*, 1977), bulbs infection in Iris (Chet *et al.*, 1983), ground nut and tomatoes (Elad *et al.*, 1982), damping-off of beans (Henis, 1984) and damping-off and stem rot of cowpea plants (Adandonon *et al.*, 2004) and collar rot of chickpea (Maurya *et al.*, 2008) caused by *S. rolfii* and increased the yield. Application of an isolate of *T. (Gliocladium) virens* in association with solarization reduced southern blight of tomatoes (Ristaino *et al.*, 1991). Mukherjee & Raghu (1997) observed that *Trichoderma* spp., were highly effective in suppressing *S. rolfii* on ginger rhizomes and on several vegetables in storage. Similarly, Chakraborty & Bhawmik (1985) found *T. viride* and *T. harzianum* highly effective in the control of sunflower collar rot caused by *S. rolfii*.

Soil population of the pathogen has a direct correlation with the intensity of disease. Shokes & Gorbet (1998) reported a positive correlation of root colonization with the population of *S. rolfii*. Similarly, Khalequzzaman (2003) observed that with a gradual increase in inoculum levels in soil, plant growth, nodulation and yield per plant reduced gradually. Chang (1994) found that the time required for the death of

*Edgeworthia papyrifera* in *S. rolfsii* infested soil was inversely proportional to number of sclerotia in soil. The present report describes the efficacy of microbial antagonists viz., *Bradyrhizobium* sp., *Rhizobium* sp., *Trichoderma harzianum*, *T. pseudokoningii*, *T. polysporum* and *T. virens* in the control of *S. rolfsii* on mungbean and sunflower and persistence of their efficacy in soil.

## Materials and Methods

A field experiment was carried out in the experimental plots of the Department of Botany, University of Karachi. Microbial antagonists viz., *Bradyrhizobium* sp., *Rhizobium* sp., *Trichoderma harzianum*, *T. pseudokoningii*, *T. polysporum* and *T. virens* multiplied on sterilized rice grains for four weeks at 25°C were applied to soil @60g in 1 m long furrow (Shahzad & Ghaffar, 1989) along with 0.5 g sclerotia of *S. rolfsii*. Plots not treated with the pathogen and biocontrol agents served as control. There were three replicates for each treatment. Mungbean and sunflower seeds were sown in the field and after 30 days growth, plants were uprooted to assess plant length, plant weight and colonization of roots by *Sclerotium rolfsii*. The roots were washed in running tap water to remove soil particles. Ten randomly selected 1cm long root pieces from each plant were surface sterilized with 1% NaOCl solution for 5 min and transferred onto potato sucrose agar plates containing Penicillin (@100,000 units L<sup>-1</sup>) and Streptomycin (@ 0.2 g L<sup>-1</sup>). After incubation for 5 days at room temperature, root colonization (RC) by *S. rolfsii* was recorded using the following formula:

$$RC\% = \frac{\text{Number of root pieces colonized}}{\text{Total number of root pieces}} \times 100$$

Data on root colonization were converted into roots colonization index (RCI) according to a 0-5 scale of Shahzad & Ghaffar (1992) where 0=0, 1=1-10, 2=11-25, 3=26-50, 4=51-75 and 5=75-100% root pieces colonized by the pathogen.

**Persistence of the effect of microbial antagonists:** After uprooting the plants in the above experiment, persistence of the effects of microbial antagonists in soil was examined by sowing the seeds of mungbean and sunflower in the same plots without any additional amendment of the microbial antagonists or the pathogen. Plant growth and root colonization by *S. rolfsii* were recorded after 30 days growth using the method described above.

## Results

Presence of *S. rolfsii* showed significant negative effect on plant growth due to severe root colonization, whereas presence of the microbial antagonists showed significant positive effect on plant growth by reducing the colonization of roots by *S. rolfsii*. Effect of the pathogen and the antagonists varied with the host. Highest root colonization by *S. rolfsii* and significant reduction in plant growth were observed in sunflower and mungbean plants

growing in soil artificially infested with sclerotia of *S. rolfsii*. The effect of *S. rolfsii* was more evident on plant weight as compared to plant length since the plants were comparatively thinner in *S. rolfsii* infested soil. Use of biocontrol agents in *S. rolfsii* infested soil showed significant reduction in RCI accompanied by increase in plant growth. *Bradyrhizobium* sp., was found most effective ( $p < 0.01$ ) followed by *Rhizobium* sp., *T. harzianum*, *T. pseudokoningii*, *T. polysporum* and *T. virens* (Figs. 1-4). Efficacy of biocontrol agents was comparatively suppressed in *S. rolfsii* infested soil as compared to non-infested soil. There was no significant difference in plant length in infested and non-infested soils where biocontrol agents were used. However, plant weights were less in infested soils as compared to non-infested soils. The reason was again the weak stems of plants in *S. rolfsii* infested soil as compared to non-infested soils.

## Persistence of the effect of microbial antagonist:

Microbial antagonists *Rhizobium*, *Bradyrhizobium* and *Trichoderma* spp., showed a greater residual effect and reduced the infection by *S. rolfsii* and enhanced the growth of mungbean and sunflower ( $p < 0.05$ ) as compared to soil that was infested with *S. rolfsii*. Soil infested with *S. rolfsii* and amended with microbial antagonists showed greater plant growth and weight as compared to control (Figs. 5, 6).




Fig. 1. Efficacy of microbial antagonists in the control of *Sclerotium rolfsii* on sunflower in microplots.

A= Control, B= *Bradyrhizobium* sp., C= *Rhizobium* sp., D= *Trichoderma harzianum*, E= *T. pseudokoningii*, F= *T. polysporum*, G= *T. virens*, Co= Control, Sr= *S. rolfsii*




Fig. 2. Efficacy of microbial antagonists in the control of *Sclerotium rolfii* on mungbean in microplots.

A= Control, B= *Bradyrhizobium* sp., C= *Rhizobium* sp., D= *Trichoderma harzianum*, E= *T. pseudokoningii*, F= *T. polysporum*, G= *T. virens*, Co= Control, Sr= *S. rolfii*

## Discussion

*Trichoderma* species are present in nearly all agricultural soils. These fungi are known to coil around the hyphae of other fungi them in a lectin-mediated reaction, and degrade their cell walls. This mycoparasitism limits growth and activity of plant pathogenic fungi. In addition to the mycoparasitism, individual strains can also produce antibiotics (Dennis & Webster, 1971). *Trichoderma* species have been used for the control of a variety of fungal pathogens like *Rhizoctonia*, *Sclerotinia sclerotiorum*, *Pythium* and *Fusarium* spp. (Harman, 1991; Lumsden & Locke, 1989; Taylor *et al.*, 1993; Lewis & Lumsden, 2001). Production of the antifungal antibiotics, gliotoxin and gliovirin, by *T. virens* has been associated with its efficacy as a biocontrol agent of most soil-borne diseases (Highley *et al.*, 1997). *T. virens* has shown promise as a preventive treatment for the control of *Rhizoctonia solani* (Howell & Stipanovic, 1995). *Trichoderma* spp., also produce organic acids, such as gluconic, citric or fumaric acids, that decrease soil pH and permit the solubilization of phosphates, micronutrients and mineral cations like iron, manganese and magnesium, useful for plant metabolism (Benitez *et al.*, 2004; Harman *et al.*, 2004). It is well documented that some strains promote plant growth, increase nutrient availability, improve crop production and enhance disease resistance (Harman *et al.*, 2004). Rhizobia have shown good potential as biological control agents against some plant pathogens. During the present studies,

use of *Rhizobium* and *Bradyrhizobium* species promoted plant growth and provided significant reduction in colonization of roots by *S. rolfii*. Blum *et al.*, (1989) reported that *Rhizobium* strains decreased the incidence of damping-off disease in Bush bean and the index of disease severity caused by *Rhizoctonia solani*. Hossain (2000) found that treatment of seeds with biofertilizer resulted in more than 85 and 73% reduction in death of plants due to infection by *Fusarium oxysporum* in lentil and chickpea, respectively. The highest reduction (69%) of fusarial foot and root rot in chickpea over untreated control was with *Rhizobium* inoculation @ 50 g/kg seed when moisten with molasses. Hossain (2000) also found that treatment of seeds with biofertilizer also showed 76 and 87% reduction in death of plants of lentil and chickpea, respectively, due to infection by *S. rolfii*. *Rhizobium* strain gave the highest promotion in plant growth. Similar results on the use of rhizobia have been reported by Hossain (2000), Kumar *et al.*, (2001) and Kibria & Hossain (2002). Solaiman (1999) found that number of total nodules per plant of chickpea was significantly increased by *Rhizobium* inoculation. Strain of *R. meliloti* are antagonistic to *F. oxysporum* (Antoun *et al.*, 1998), and rhizobia antagonistic to *F. solani* isolated from commercial snap bean, appeared to have a good potential for controlling *Fusarium* rot (Buonassisi *et al.*, 1986). Ehtesham-ul-Haque & Ghaffar (1993) observed under field conditions that *R. meliloti*, *R. leguminosarum* and *B. japonicum* used either as soil drench or seed dressing reduced infection of roots by *Macrophomina phaseolina*, *Rhizoctonia solani* and *Fusarium* spp., in leguminous as well as non-leguminous plants. *Rhizobium leguminosarum* significantly increased the number and growth of pea seedlings (Bardin *et al.*, 2004). Attachment of rhizobia to roots of non-leguminous crops has been described by Terouchi & Syono (1990) that can explain the efficacy of rhizobia on non-leguminous crops.

*Bradyrhizobium* and *Rhizobium* share characteristic with plant growth promoting rhizobacteria (PGPR). Like other PGPR, these nodule inducing bacteria colonize the roots of non-legumes, produce phytohormones, siderophores and HCN and exhibit antagonistic effects towards plant pathogenic fungi (Antoun *et al.*, 1998). The increment in growth parameters in response to rhizobial inoculation endorsed the fact that the test strains were having one or more growth promoting mechanisms including mobilization and efficient uptake of nutrients (Biswas *et al.*, 2000 a, b), enhancement in stress resistance (Alami *et al.*, 2000), solubilization of insoluble phosphates (Alikhani *et al.*, 2006), induction of systemic disease resistance (Reitz *et al.*, 2000), inhibition of fungal growth (Nautiyal, 1997), production of phytohormones (Dakora, 2003), vitamins (Dobbelaere *et al.*, 2003) and siderophores (Neiland & Leong, 1986). Biswas *et al.*, (2000b) reported 16% increase in number of panicles per plant of rice and suggested that the improvement was due to increase availability of nutrients and phytohormones like indole acetic acid and ethylene. Similarly, Chi *et al.*, (2005) observed more than 23% increase in plant height of rice over uninoculated control and argued indole acetic acid and gibberellins production as the key mechanism for that improvement. Our results corroborate well with those of Gaur *et al.*, (1980) who found that rhizobia and bradyrhizobia, significantly increased the weight of shoot, number of pods, nodule volume and the yield of the following green gram (*Vigna radiata* (L.) Wilczek) and groundnut (*Arachis hypogaea* L.) crops. Pena-Cabriaes & Alexander (1983) found that strain of rhizobia and




Fig. 3. Effect of microbial antagonists on growth of sunflower plants.  
A= Control, B= Pathogen alone, C= Antagonists alone, D= Antagonist and pathogen together.




Fig. 4. Effect of microbial antagonists on growth of mungbean plants.  
A= Control, B= Pathogen alone, C= Antagonists alone, D= Antagonist and pathogen together




Fig. 5. Persistence of the efficacy of microbial antagonists in the control of *Sclerotium rolfsii* on sunflower in microplots.

A= Control, B= *Bradyrhizobium* sp., C= *Rhizobium* sp., D= *Trichoderma harzianum*, E= *T. pseudokoningii*, F= *T. polysporum*, G= *T. virens*, Co= Control, Sr= *S. rolfsii*

bradyrhizobia grew readily in the presence of germinating seeds and developing root systems of soybean (*Glycine max* (L.) Merr.), red clover (*Trifolium pratense* L.), kidney beans (*Phaseolus vulgaris* L.), cowpeas (*Vigna unguiculata* L.), oats, wheat and corn. Reitz *et al.*, (2000) have reported the induction of systemic resistance to the cyst nematode *Globodera pallida* in potato by *Rhizobium etli* strain G12. The beneficial effect of *Rhizobium* and *Bradyrhizobium* in legumes in terms of biological nitrogen fixation has been a main focus in the recent past. The results of the present study would suggest that use of *Trichoderma*, *Rhizobium* and *Bradyrhizobium* species can provide protection against *S. rolfsii* infection resulting in increased crop growth and productivity. The ability of these biocontrol agents to persist in soil can also provide protection in the next crop as well.

## References

- Adandonon, A. 2000. *Damping-off of cowpea in Benin and South Africa*. M.Sc. Thesis. University of Pretoria, Pretoria, South Africa.
- Adandonon, A., Aveling, T.A.S. and M. Tamo. 2004. Occurrence and distribution of cowpea damping-off and stem rot and associated fungi in Benin. *The Journal of Agricultural Science*, 142(5): 561-566.




Fig. 6. Persistence of the efficacy of microbial antagonists in the control of *Sclerotium rolfsii* on mungbean in microplots.

A= Control, B= *Bradyrhizobium* sp., C= *Rhizobium* sp., D= *Trichoderma harzianum*, E= *T. pseudokoningii*, F= *T. polysporum*, G= *T. virens*, Co= Control, Sr= *S. rolfsii*

- Agrawal, S.C., M.N. Khare and P.S. Agrawal. 1977. Biological control of *Sclerotium rolfsii* causing collar rot of lentil. *Indian Phytopathology*, 30: 176-179.
- Ahmed, Y., M.S. Mirza and M. Aslam. 1984. *Sclerotium rolfsii* on maize. *FAO Plant Prot. Bull.*, 32: 147.
- Alami, Y., W.A.C. Achouak and T. Heulin. 2000. Rhizosphere soil aggregation and plant growth promotion of sunflowers by an exopolysaccharide producing *Rhizobium* sp., strain isolated from sunflower roots. *Applied & Environmental Microbiology*, 66: 3393-3398.
- Alikhani, H.A., N. Saleh-Rastin and H. Antoun. 2006. Phosphate solubilization activity of rhizobia native to Iranian soils. *Plant & Soil*, 287: 35-41.
- Antoun, H., C.J. Beauchamp, N. Goussard, R. Chabot and R. Lalande. 1998. Potential use of *Rhizobium* and *Bradyrhizobium* growth promoting rhizobacteria with nonleguminous radishes (*Raphanus sativus* L.). *Plant & Soil*, 204: 57-67.
- Atlas, R. and R. Bartha. 1998. *Microbial Ecology. Fundamental and Applications*. Benjamin/Cummings Sci. Pub., Menlo Park, CA
- Aycock, R. 1966. Stem rot and other diseases caused by *Sclerotium rolfsii*. *NC. Agric. Exp. Sta. Tech. Bull.*, No. 174. 202pp.
- Bardin, S.D., H.C. Huang, J. Pinto, E.J. Amundsen and R.S. Erickson. 2004. Biological control of *Pythium* damping-off of pea and sugar beet by *Rhizobium leguminosarum* pv *viceae*. *Can. J. Bot.*, 82: 291-296.

- Benitez, T., A.M. Rincon and A.C. Condon. 2004. Biocontrol mechanism of *Trichoderma* strains. *International Microbiology*, 7: 249-260.
- Biswas, J.C., J.K. Ladha and F.B. Dazzo. 2000a. Rhizobia inoculation improves nutrient uptake and growth of lowland rice. *Soil Science Society of America Journal*, 92: 880-886.
- Biswas, J.C., J.K. Ladha, F.B. Dazzo, Y.G. Yanni and B.G. Rolfe. 2000b. Rhizobial inoculation influences seedling vigor and yield of rice. *Agronomy Journal*, 92: 880-886.
- Blum, L.K., S.D. Frey and G. Soto. 1989. Effect of fluorescent-pigment producing *Rhizobium* on the severity of *Rhizoctonia solani* seed and root rot of *Phaseolus vulgaris*. A paper presented at a symposium held at the Beltsville Agricultural Research Center (BARC), 8-11 May, 1989.
- Buonassisi, A.J., R.J. Copeman, H.S. Pepin and G.W. Eaton. 1986. Effect of *Rhizobium* spp. on *Fusarium solani* f.sp. *phaseoli*. *Can. J. Plant Pathol.*, 8: 140-146.
- Chakraborty, S. and T.P. Bhawmik. 1985. Chemical and biological control of *Sclerotium rolfsii*. *Pesticides*, 19(2): 31-33.
- Chang, T. 1994. Pathogenicity of *Sclerotium rolfsii* on *Edgeworthia papyrifera* and sclerotial survival. *Bull Taiwan For. Res. Inst. New Series*, 9: 191-196.
- Chet, I., I. Elad, A. Kalfon, Y. Hadar and J. Katan. 1983. Integrated control of soilborne and bulb borne pathogens in Iris. *Phytoparasitica*, 10: 229.
- Chi, F., H.S. Shen, H.P. Cheng, Y.X. Jing, Y.G. Yanni and F.B. Dazzo. 2005. Ascending migration of endophytic rhizobia, from roots to leaves, inside rice plants and assessment of benefits to rice growth physiology. *Applied & Environmental Microbiology*, 71: 7271-7278.
- Ciancio, A. and K.G. Mukerji. 2007. *General concepts in integrated pest and disease management*. Springer. 359 pp.
- Ciccarese, F., S. Frisullo, M. Amenduni and M. Cirulli. 1992. Biological control of *Sclerotium rolfsii* root rot of sugar beet with *Trichoderma harzianum*. pp. 243-248. In: (Eds.): E.C. Tjamos, G.C. Papavizas and R.J. Cook. Plenum Press New York.
- Cilliers A.J., L. Herselman and Z.A. Pretorius. 2000. Genetic variability within and among mycelial compatibility groups of *Sclerotium rolfsii* in South Africa. *Phytopathology*, 90: 1026-1031.
- Dakora, F.D. 2003. Defining new roles for plant and rhizobial molecules in sole and mixed plant cultures involving symbiotic legumes. *New Phytol.* 158: 39-49.
- Dennis, C. and J. Webster. 1971. Antagonistic properties of species-groups of *Trichoderma*: II. Production of volatile antibiotics. *Trans. Brit. Mycol. Soc.*, 57(1): 41-48.
- Dobbelaere, S., J. Vanderleyden and Y. Okon. 2003. Plant growth promoting effects of diazotrophs in the rhizosphere. *Plant & Soil*, 22: 107-149.
- Domsch, K.H., W. Gams and T.H. Anderson. 1980. *Compendium of Soil Fungi*. Academic Press, London, New York.
- Ehteshamul-Haque, S. and A. Ghaffar. 1993. Use of *Rhizobia* in the control of root rot diseases of sunflower, okra, soy bean and mung bean. *J. Phytopathology*, 138: 157-163.
- Elad, Y., I. Chet and J. Katan. 1980. *Trichoderma harzianum*: A biocontrol agent effective against *Sclerotium rolfsii* and *Rhizoctonia solani*. *Phytopathology*, 70: 119-121.
- Elad, Y., Y. Hadar, I. Chet and Y. Henis. 1982. Prevention with *Trichoderma harzianum* Rifai aggr., of reinfestation by *Sclerotium rolfsii* Sacc., and *Rhizoctonia solani* Kuhn of soil fumigated with methyl bromide and improvement in disease control in tomato and peanuts. *Crop Prot.*, 1: 199-211.
- Farr, D.F., G.F. Bills, G.P. Chamuris and A.Y. Rossman. 1989. *Fungi on plant and plant product in the United States*. American Phytopathological Society, St. Paul.
- Gaur, Y.D., A.N. Sen and N.S. Subba Rao. 1980. Improved legume-rhizobium symbiosis by inoculating preceding cereal crop with Rhizobium. *Plant & Soil*, 54: 313-316.
- Harman, G.E. 1991. Seed treatments for biological control of plant disease. *Crop Protection*, 10(3): 166-171.
- Harman, G. E., C.R. Howell, A. Viterbo, I. Chet and M. Lorito. 2004. *Trichoderma* species—Opportunistic, avirulent plant symbionts. *Nat. Rev. Microbiol.*, 2: 43-56.
- Henis, Y. 1984. Interactions between *Sclerotium rolfsii* and *Trichoderma* spp., relationship between antagonism and disease control. *Soil Biology & Biochemistry*, 16(4): 391-395.
- Henis, Y., P.B. Adams, J.A. Lewis and G.C. Papavizas. 1983. Penetration of sclerotia of *Sclerotium rolfsii* by *Trichoderma* spp. *Phytopathology*, 73: 1043-1046.
- Highley, T.L., H.S.A. Padmanabha and C. R. Howell. 1997. Control of wood decay by *Trichoderma* (*Gliocladium*) *virens*. II. Antibiosis. *Mat. und Org.*, 31(3): 157-166.
- Hossain, I. 2000. Biocontrol of *Fusarium oxysporum* and *Sclerotium rolfsii* infection in lentil, chickpea and mungbean. *BAU Res. Prog.*, 11: 61.
- Howell, C.R. and R.D. Stipanovic. 1995. Mechanisms in the biocontrol of *Rhizoctonia solani*-induced cotton seedling disease by *Gliocladium virens*: Antibiosis. *Phytopathology*, 85: 469-472.
- Iqbal, S.M., A. Bakhsh, S. Hussain and B.A. Malik. 1995. Microbial antagonism against *Sclerotium rolfsii*, the cause of collar rot of lentil. *Lens-Newsletter*, 22: 48-49.
- Jahangir, A.M., M.K. Mohibullah, M. Ayub and M. Khan. 1995. Detection of fungi causing root rot of apple nurseries in Swat. *Sarhad J. Agri.*, 11: 754-748.
- Khalequzzaman, K.M. 2003. Effect of inocula levels of *Meloidogyne javanica* and *Sclerotium rolfsii* on the plant growth, yield and galling incidence of soybean. *Pak. J. Pl. Pathol.*, 2: 56-64.
- Kibria, M.G. and I. Hossain. 2002. Effect of biofertilizer and *Rhizobium* on foot and root rot disease and seed yield of mungbean. *Bangladesh J. Seed Sci. & Tech.*, 6(1&2): 1-6.
- Kossou, D.K., G. Gbèhounou, A. Ahanchédé, B. Ahohuendo, Y. Bouraïma and A. van Huis. 2001. Indigenous Cowpea Production and Protection Practices in Benin. *Insect Sci. Applic.*, 21(2): 123-132.
- Kumar, R., P.P. Gupta and B.L. Jalali. 2001. Impact of VA-mycorrhiza, *Azotobacter* and *Rhizobium* on growth and nutrition of cowpea. *Plant the field on germination of sclerotia. Plant Disease*, 80: 655-659.
- Lewis, J.A. and R.D. Lumsden. 2001. Biocontrol of damping-off of greenhouse-grown crops caused by *Rhizoctonia solani* with a formulation of *Trichoderma* spp. *Crop Protection*, 20(1): 49-56.
- Lumsden, R.D. and J.C. Locke. 1989. Biological control of *Pythium ultimum* and *Rhizoctonia solani* damping-off with *Gliocladium virens* in soil-less mix. *Phytopathology*, 79: 361-366.
- Maurya, S., R. Singh, D.P. Singh, H.B. Singh, U.P. Singh and J.S. Srivastava. 2008. Management of collar rot of chickpea (*Cicer arietinum*) by *Trichoderma harzianum* and plant growth promoting rhizobacteria. *J. Plant Protect. Res.*, 48(3): 347-354.
- Mukherjee, P.K. and K. Raghu. 1997. *Trichoderma* species as a microbial suppressive agent of *Sclerotium rolfsii* on vegetables. *World J. Microbiol. Biotechnol.*, 13: 497-499.
- Nautiyal, C.S. 1997. Rhizosphere competence of *Pseudomonas* sp. NBRI9926 and *Rhizobium* sp. NBRI9513 involved in the suppression of chickpea (*Cicer arietinum* L.) pathogenic fungi. *FEMS Microbiology Ecology*, 23(2): 145-158.
- Neiland, J.B. and S.A. Leong. 1986. Siderophores in relation to plant growth and disease. *Annual Review in Plant Physiology*, 37: 187-208.

- Papavizas, G.C. and J.A. Lewis. 1989. Effect of *Gliocladium* and *Trichoderma* on damping-off and blight of snap bean caused by *Sclerotium rolfsii* in the greenhouse. *Plant Pathol.*, 38: 277-286.
- Papavizas, G.C. and R.D. Lumsden. 1980. Biological control of soilborne fungal propagules. *Ann. Rev. Phytopathol.*, 18: 389-413.
- Pena-Cabriaes J.J. and M. Alexander. 1983. Growth of *Rhizobium* in unamended soil. *Soil Sci. Soc. Am. J.*, 47: 81-84.
- Punja, Z.K. 1985. The biology, ecology and control of *Sclerotium rolfsii*. *Annual review of Phytopathology*, 23: 97-127.
- Punja, Z.K. 1988. *Sclerotium (Athelia) rolfsii*, a pathogen of many plant species. pp. 523-534. Vol. 6. In: *Advances in Plant Pathology*. (Ed.): G.S. Sidhu. Academic Press, London.
- Reitz, M., K. Rudolph, I. Schroder, S. Hoffmann-Hergarten, J. Hallmann and R.A. Sikora. 2000. Lipopolysaccharides of *Rhizobium etli* strain G12 act in potato roots as an inducing agent of systemic resistance to infection by the cyst nematode *Globodera pallida*. *Appl. Environ. Microbiol.*, 66: 3515-3518.
- Ristaino, J.B., K.B. Perry and R.D. Lumsden. 1991. Effect of soil solarization and *Gliocladium virens* on sclerotia of *Sclerotium rolfsii*, soil microbiota, and the incidence of southern blight in tomato. *Phytopathology*, 81: 1117-1124.
- Rolfs, P.H. 1892. Tomato blight, some hints. Bulletin of Florida Agricultural Experimental Station. p.18.
- Ruqia, B. 2001. *Studies on seed-borne mycoflora of sugarbeet (Beta vulgaris)*. M.Sc. Thesis, Department of Botany, University of Karachi, Karachi. 30 pp.
- Shahzad, S. and A. Ghaffar. 1989. Use of *Paecilomyces lilacinus* in the control of root rot and root knot disease complex of okra and mungbean. *Pak. J. Nematol.*, 7: 47-53.
- Shahzad, S. and A. Ghaffar. 1992. Root rot and root knot disease complex of mungbean and its biological control. pp. 349-256. In: *Status of Plant Pathology in Pakistan. Proc. National Symp.*, Ghaffar, A. & Shahzad, S. (Eds.). Department of Botany, University of Karachi.
- Shahzad, S. and A. Ghaffar. 1995. New records of soilborne root infecting fungi in Pakistan. *Pak. J. Bot.*, 17: 209-216.
- Shokes, F. and D. Gorbet. 1998. Crop losses due to stem rot of groundnut in commercial cultivars and partially resistant breeding lines. *ICPP98 Paper No. 3.4.5*.
- Solaiman, A.R.M. 1999. Influence of *Rhizobium* inoculant, nitrogen and boron on nodulation, dry weight and grain yield of chickpea. *Annals of Bangladesh Agric.*, 9: 75-84.
- Taylor, G.A., G.E. Harman and A.P. Nielson. 1993. Biological seed treatments using *Trichoderma harzianum* for horticultural crops. In: *New chemical and biological treatments for horticultural seeds. Hort. Technology*, 4: 105-109.
- Terouchi, N. and Syōno, K. 1990. *Rhizobium* attachment and curling in asparagus, rice and oat plants. *Plant & Cell Physiology*, 31(1): 119-127.
- Tjamos, E.C., G.C. Papavizas and R.J. Cook. 1992. *Biological control of plant diseases. Progress and challenges for the future*. Plenum Press, New York, NY. pp. 125-132.
- Yorinori, J.T. 1994. Fungal Diseases. pp. 37-60. In: *Tropical soybean: improvement and production*. Embrapa (ed.). FAO.

(Received for publication 3 October 2009)