

## EFFECT OF THIDIAZURON (TDZ) ON IN VITRO MICROPROPAGATION OF *SOLANUM TUBEROSUM* L. CVS. DESIREE AND CARDINAL

ZAHOOR AHMAD SAJID AND FAHEEM AFTAB\*

Department of Botany,  
University of the Punjab, Q. A. Campus Lahore-54590, Pakistan.

### Abstract

Thidiazuron (1-phenyl-3- (1,2,3- Thiadiazol-5-yl) urea; TDZ) is one of the several substituted ureas that have been investigated recently for their cytokinin-like activity. TDZ is known to be more active than zeatin for stimulating the growth when added to a tissue culture medium at a low concentration. In this study, effect of TDZ on several *In vitro* growth parameters in *Solanum tuberosum* cvs. Desiree and Cardinal were observed. Shoot apices (1.0cm each) from both the cultivars were separately inoculated on full strength MS basal media as well as on MS full strength supplemented with different concentrations of TDZ ( $10^{-10}$ ,  $10^{-9}$  or  $10^{-8}$  M) thus forming four combinations. Results were recorded for shoot length, shoot number, root length, root number, number of nodes, fresh and dry weight of plants after 30 days of explant inoculation. MS full strength medium was found to be the best for *In vitro* micropropagation. However, TDZ did have an influence on the studied growth parameters, which have been discussed and interpreted herein.

### Introduction

TDZ a substituted phenylurea (N-phenyl-N-1,2,3-thiadiazol-5-ylurea) is used as a synthetic herbicide and a plant growth regulator to stimulate high rate of axillary shoot proliferation in many woody plant species (Malik & Saxena, 1992). It has been considered to be more potent than most of the commonly used cytokinins (Huetteman & Preece, 1993). Thidiazuron releases the lateral bud dormancy and stimulates shoot formation in wide variety of plant species (Fiola *et al.*, 1990; Malik & Saxena, 1992). According to Capelle *et al.*, (1983), TDZ directly promotes growth due to its own biological activities in a fashion similar to that of an N- substituted cytokinin or it may induce the synthesis and accumulation of an endogenous cytokinin. In woody plant species, low levels of TDZ induce the axillary shoot proliferation but higher levels may inhibit it. Higher levels, on the other hand, promote callus and somatic embryo formation (Huetteman & Preece, 1993).

Potato is one of the most important widely-grown crops and is an integral part of diet in the entire world. It produces more protein (524 kg/ha) as compared to wheat (254kg/ha). It also supplies at least 12 essential minerals including Vitamin C (Irfan, 1992). Apart from woody plant species, TDZ has shown promise in many other plants belonging to diverse groups or families such as *Camellia sinensis* (Mondal *et al.*, 1998), *Hordeum vulgare* (Ganesan *et al.*, 2003), *Oryza sativa* (Gairi & Rashid, 2004), *Hyoscyamus niger* (Uranbey 2005). To our knowledge, it has never been tested for *In vitro* clonal propagation of potato. It is important that the effect of TDZ be tested on important crops like potato. The goal of this research work was to study the effect of TDZ on growth of *Solanum tuberosum* L., using shoot apices as explant from *In vitro* raised plants.

\*E-mail: faheem@botany.pu.edu.pk

## Materials and Methods

Apical shoot explants of *Solanum tuberosum* L. cvs. Desiree and Cardinal were grown on MS (Murashige & Skoog, 1962) full strength or MS medium containing three different concentrations of TDZ ( $10^{-8}$ ,  $10^{-9}$  or  $10^{-10}$  M). All media were solidified with 0.7% Agar- agar (Merck) and adjusted to pH 5.8. Apical shoot explants (ca. 1.0 cm) from both the cultivars were grown in each medium. Ten culture vessels were used for each treatment per experiment and each experiment was repeated thrice for the two varieties. The cultures were incubated under 16-h photoperiod ( $35 \mu\text{mol m}^{-2} \text{ s}^{-1}$  from cool fluorescent lights) at  $25 \pm 2^\circ\text{C}$ .

Results were recorded for shoot length, shoot number, root length, root number, number of nodes, fresh and dry weight of plants after 30 days of explant inoculation. For this purpose, the plants were harvested and shoot length was recorded with the help of a suitable scale from the top of the medium to the tip of shoot minus 1.0 cm (size of the original explant). The root length was measured from the tip of the root up to the basal end of the shoot. Fresh weights of the plantlets were determined. Afterwards, the plant material was placed in small brown envelopes and dried in hot air oven at  $60^\circ\text{C}$  for up to 48 hours to estimate dry weight.

## Results and Discussion

### *In vitro* establishment of potato shoot apices

**a. Shoot and root length:** The effect of TDZ, a substituted phenyl-urea as a plant growth regulator has been reported for many plant species including several recalcitrant woody species like *Quercus robur* L., (Chalupa, 1988) and *Pinus strobes* L., (Pijut *et al.*, 1991). In potato (cv. Cardinal), highest shoot length (5.74 cm) and root length (5.34 cm) was observed in MS full strength medium after 30 days of initial culture (Table 1). This was followed by MS + TDZ ( $10^{-10}$ ) where the shoot and root length was 5.51 cm and 5.24 cm, respectively. Lowest shoot and root length (5.30 and 4.08 cm respectively) was observed at  $10^{-8}$  M TDZ. In cv. Desiree, highest shoot and root length was 5.48 cm and 5.32 cm respectively on MS full strength medium (Table 2). A somewhat similar pattern for shoot and root data was observed since the lowest values for the two parameters were observed on MS medium supplemented with  $10^{-8}$  M TDZ.

It is evident from these results that TDZ does have an influence on micro propagation of potato even at very low concentrations as the second highest shoot and root length was recorded using MS medium supplemented with  $10^{-10}$  TDZ.

**b. Number of shoots, roots and nodes:** Maximum number of shoots (2.66 and 2.96) was obtained on MS + TDZ ( $10^{-8}$ ) in cvs. Cardinal and Desiree, respectively (Tables 1 and 2). It might be due to the high concentration of TDZ as it modifies the endogenous cytokinin metabolism (Capelle *et al.*, 1983; Hare & Van Staden, 1994; Murthy *et al.*, 1995; Hutchinson & Sexena, 1996). Alternatively, it has been suggested that TDZ may mimic an auxin response (Visser *et al.*, 1992) or modifies endogenous auxin metabolism (Murthy *et al.*, 1995; Hutchinson *et al.*, 1996). However, highest number of roots (12.60 and 14.90) and nodes (7.90 and 7.20) was observed on MS full strength medium in the two cultivars respectively. The lowest number of shoot (2.0 and 1.5), root (7.3 and 4.9) and node (6.3 and 6.3 respectively) was obtained on MS + TDZ  $10^{-9}$  in both the cultivars.



**c. Fresh and dry weight of plantlets:** The maximum fresh and dry weight of the plantlets (0.543 g and 0.0524 g) in cv. Cardinal was obtained on MS medium containing  $10^{-9}$  M TDZ. In Desiree, the highest fresh and dry weights (1.0560 and 0.0965 g, respectively) were observed on MS medium containing  $10^{-10}$  M TDZ (Tables 1 and 2). Although the highest shoot and root length (in both the cultivars) was obtained on MS full strength medium but highest fresh and dry weights in Desiree and Cardinal were observed using MS medium with either  $10^{-10}$  or  $10^{-9}$  TDZ, respectively.

The results from the present work demonstrate the possibility to micropropagate potato using TDZ that to our knowledge has not been tested before in potato for the purpose of *In vitro* clonal propagation. Its superiority over general basal MS medium, however, could not be established in this study. It also demonstrates that TDZ might be used at very low concentration (less than a nM level) for potato micropropagation as compared to other cytokinins (such as BAP) that are usually used at relatively higher ( $\mu$ M) levels (Rida *et al.*, 2001; Fengyen & Han 2002). Although TDZ is relatively costly, a very low concentration of TDZ used in our studies circumvents its price consideration and seems to be quite cost-effective. Moreover, its influence on all the growth parameters under study necessitates further work using TDZ as a growth regulator to better understand its role in potato tissue culture.

## Reference

- Capelle, S.C., D.W.S. Mok, S.C. Kirchner and M.C. Mok. 1983. Effects of thiadiazuron on cytokinin autonomy and metabolism of  $N^6$ - (2- isopentyl) [ $8-^{14}C$ ] adenosine in callus tissue of *Phaseolus lunatus* L. *Plant Physiol.*, 73: 796- 802.
- Chalupa, V. 1988. Large-scale micropropagation of *Quercus robur* L., using adenine type cytokinin and thidiazuron to stimulate shoot proliferation. *Biol. Plant (praha)*, 30: 414-421.
- Fengyen, L. and L. Han. 2002. Effect of exogenous hormones on micropropagation of *In vitro* virus free potato plantlets. *Chines Potato J.*, 16: 214-216.
- Fiola, J.A., M.A. Hassan, H.J. Swartz, R.H. Bors and R. McNicols. 1990. Effect of thidiazuron, light influence rates and kanamycin on *In vitro* shoot organogenesis from excised Rubus cotyledons and leaves. *Plant. Cell. Tiss. Org. Cult.*, 20: 223-228.
- Gairi, A. and A. Rashid. 2004. TDZ induced somatic embryogenic in non-responsive caryopsis of rice using shot treatment with 2,4-D. *Plant. Cell. Tiss. Org. Cult.*, 76: 29-33.
- Ganesan, S., B. Monica, L.H. Bryan, G.R. Brian, J.S. Graham and N.C. Ravindra. 2003. Production of multiple shoots from thidiazuron-treated mature embryos and leaf-base/apical meristem of barley (*Hordeum vulgare*). *Plant Cell. Tiss. Org. Cult.*, 73: 57-64.
- Hare, P.D. and J. Van Staden. 1994. Inhibitory effect of TDZ on the activity of cytokinin oxidase isolated from soybean callus. *Plant Cell. Physiol.*, 35: 1121-1125.
- Huetteman, A. and E.J. Preece. 1993. Thidiazuron: a potent cytokinin for woody plant tissue culture. *Plant Cell. Tiss. Org. Cult.*, 33: 105-119.
- Hutchinson, M.I. and P.K. Saxena. 1996. Role of purine metabolism in TDZ induced somatic embryogenesis of geranium (*Pelargonium hortorum* L. Bailey). *J. Plant Physiol.*, 149: 573-579.
- Hutchinson, M.I., S. Krishna Raj and P.K. Saxena. 1996. Morphological and physiological changes during thiadiazuron induced somatic embryogenesis in geranium (*Pelargonium hortorum* L.) Hypocotyl culture. *Int. J. Plant. Sci.*, 157: 110-117.
- Irfan, A. 1992. *Proceedings of the workshop on Seed Potato Health*. Seed Potato Health Laboratory Project, Sahiwal. Federal Seed Certification Department. Ministry of Food, Agriculture and Cooperation, pp. 1-2.
- Malik, K.A. and P.K. Saxena. 1992. Regeneration in *Phaseolus vulgaris* L., high frequency induction of direct shoot formation in intact seedlings by  $N^6$ - benzylaminopurine and thidiazuron. *Planta*, 186: 384-389.

- Mondal, T.K., B. Amita, S. Anil and S.A. Paramvir. 1998. Micropropagation of tea (*Camellia sinensis*) using thidiazuron. *Plant Cell. Tiss. Org. Cult.*, 26: 57-61.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.*, 15: 473- 497.
- Murthy, B.N.S., S.J. Murch and P.K. Saxena. 1995. Thidiazuron induced somatic embryogenesis in intact seedling of peanut (*Arachis hypogaea* L.) Endogenous growth regulator level and significance of cotyledons. *Phys. Plant.*, 94: 268-276.
- Pijut, P.M., C.H. Michler and T.M. Voelker. 1991. Effect embryo explant orientation, thidiazuron, and agar on eastern white pine (*Pinus strobus* L.) adventitious shoot initiation. *Proc. intl. Symposium on application of biotechnology to tree culture, protection and utilization.* Columbus OH (pp126) (Abstr).
- Rida, A., A.M. Shibli, A. Suwan and M. Ajlouni. 2001. *In vitro* multiplication of virus free Spunta potato. *Pak. J. Bot.*, 33(1): 35: 41.
- Uranbey, S. 2005. Thidiazuron induced adventitious shoot regeneration in *Hyoscyamus niger*. *Bilogia. Plantarum*, 49(3): 427-430.
- Visser, C., J.A. Qureshi, R. Gill, K.A. Malik and P.K. Saxena. 1992. Morpho-regulatory role of thidiazuron: Substitution of auxin and cytokinin requirement for the induction of somatic embryogenesis in geranium hypocotyls culture. *Plant Physiol.*, 99: 1704-1707.

(Received for publication 24 August 2006)