

## SEED-BORNE MYCOFLORA OF WHEAT, SORGHUM AND BARLEY

FAKHRUNNISA, M.H. HASHMI\* AND A. GHAFFAR\*

Department of Botany,  
Khatoon-e-Pakistan Government Degree College for Women,  
Karachi, Pakistan

### Abstract

Standard blotter and Deep Freezing methods were used to study the seed-borne mycoflora of 19 samples of wheat, 27 samples of sorghum and 14 samples of barley. A significant contamination with fungal genera was detected in analyzed samples. Fungi most frequently isolated and identified were *Absidia* sp., *Alternaria alternata*, *Aspergillus* sp., *A. candidus*, *A. flavus*, *A. niger*, *A. sulphureus*, *Cephalosporium* sp., *Chaetomium globosum*, *Cladosporium herbarum*, *Curvularia lunata*, *Drechslera dematioidea*, *D. halodes*, *D. hawaiiensis*, *D. tetramera*, *Fusarium moniliforme*, *F. oxysporum*, *F. pallidoroseum*, *F. subglutinans*, *Nigrospora oryzae*, *Penicillium* spp., *Piptocephalis* sp., *Rhizoctonia solani*, *Rhizopus* sp., *Stemphylium* sp., *Syncephalastrum racemosum*, *Trichoderma hamatum*, *Trichothecium roseum* and *Ulocladium* sp. This is the first report of *Chaetomium globosum* and *D. hawaiiensis* on wheat, *A. sulphureus*, *Fusarium subglutinans*, *Nigrospora oryzae*, *Piptocephalis* sp., *Syncephalastrum racemosum* and *Trichoderma hamatum* on sorghum and *A. niger*, *Cephalosporium* sp., *Cladosporium herbarum*, *Drechslera dematioidea*, *D. tetramera*, *Trichothecium roseum*, *Stemphylium* sp., and *Ulocladium* sp., are new records on barley. There does not appear to be any previous report of *Absidia* sp., *Aspergillus sulphureus*, *Fusarium subglutinans* and *Rhizoctonia solani* on wheat in Pakistan. Deep freezing method showed better results for isolation of *Alternaria alternata*, *Cladosporium herbarum*, *Drechslera* spp., and *Fusarium* spp.

### Introduction

Seeds are regarded as highly effective means for transporting plant pathogens over long distances. Numerous examples exist in agriculture literature for the international spread of plant diseases as a result of the importation of seeds that were infected or contaminated with pathogens (Agarwal & Sinclair, 1996).

Seed-borne diseases have been found to affect the growth and productivity of crop plants (Kubiak & Korbas, 1999; Weber *et al.*, 2001; Dawson & Bateman, 2001). A seed-borne pathogen present externally or internally or associated with the seed as contaminant, may cause seed abortion, seed rot, seed necrosis, reduction or elimination of germination capacity as well as seedling damage resulting in development of disease at later stages of plant growth by systemic or local infection (Khanzada *et al.*, 2002; Bateman & Kwasna, 1999).

Wheat is one of the main staple food of man and is grown in almost all the temperate and subtropical regions of the world. Seed-borne mycoflora of wheat reported recently included *Alternaria alternata*, *Drechslera sorokiniana*, *Fusarium moniliforme*, *F. avenaceum*, *F. graminearum*, *F. nivale*, *F. culmorum*, *F. equiseti*, *F. sporotrichioides*, *Cladosporium herbarum*, *Stemphylium botryosum* (Nirenberg *et al.*, 1994; Glazek, 1997; Mirza & Qureshi, 1978).

Seed-borne mycoflora of sorghum reported from different parts of the world include *Alternaria alternata*, *Aspergillus flavus*, *A. fumigatus*, *A. niger*, *Cladosporium* sp., *Fusarium moniliforme*, *F. oxysporum*, *F. pallidoroseum*, *Drechslera tetramera*, *Nigrospora* sp., *Phoma* sp., and *Rhizopus* sp., (Abdullah and Kadhum, 1987; Ahmed *et al.*, 1992).

\*Department of Botany, University of Karachi, Karachi, Pakistan.

Some of the seed-borne mycoflora of barley include *Alternaria alternata*, *Aspergillus candidus*, *A. flavus*, *Curvularia lunata*, *Drechslera halodes*, *Fusarium moniliforme*, *F. pallidoroseum*, *F. solani* and *Ulocladium* sp., (Richardson, 1983; Ahmed et al., 1997).

Besides, the mold fungi which grow on the seed substratum produce mycotoxins which are hazardous to man and animals (Halt, 1994). Studies were carried out to study the composition of seed-borne mycoflora occurring in wheat, sorghum and barley grains which are the main crops grown in Pakistan.

### Materials and Methods

Seed samples of *Triticum aestivum* (19 samples), *Sorghum vulgare* (27 samples) and *Hordeum vulgare* (14 samples) collected from different localities of Pakistan viz., Karachi, Lahore, Islamabad and Faisalabad were used for the isolation and detection of seed-borne fungi. Blotter Method recommended by International Seed Testing Association (Anon., 1966) and Deep Freezing Method (Limonard, 1968) were used for the isolation of fungi.

**Blotter Method:** Four hundred seeds of each sample were plated on three layers of moistened blotters placed in 90 mm diam., Petri plates @ 25 seeds/plate. The plates were incubated for 7 days at  $20^{\circ}\text{C} \pm 1^{\circ}\text{C}$ . Then were examined under the microscope for seed-borne mycoflora.

**Deep Freezing Method:** Twenty five seeds per plate were plated on three layers of moistened blotters. The seeds were incubated for one day at  $22^{\circ}\text{C}$  followed by 24 hours of freezing at  $-20^{\circ}\text{C}$ . The plates were then placed for 4-5 days at  $22^{\circ}\text{C} \pm 1^{\circ}\text{C}$ .

After incubation the growth characters as well as percentage of infection were recorded. The isolated *Fusarium* spp., were maintained on Spezieller Nahrstofffarmer agar (SNA) medium (Nirenberg, 1976; Hashmi & Thrane, 1990) whereas other fungi were maintained on PDA. The fungi were identified after reference to Barnett & Hunter (1972), Booth (1971), Nelson et al., (1983).

### Results and Discussion

The occurrence of fungi most frequently encountered is recorded in terms of mean value with standard error (Table 1).

**a. Wheat:** From 19 samples of wheat, 12 genera and 21 species of fungi viz., *Absidia* sp., *Alternaria alternata*, *Aspergillus* sp., *A. candidus*, *A. flavus*, *A. niger*, *A. sulphureus*, *Cephalosporium* sp., *Chaetomium globosum*, *Cladosporium herbarum*, *Curvularia lunata*, *Drechslera halodes*, *D. hawaiiensis*, *D. tetramera*, *Fusarium moniliforme*, *F. oxysporum*, *F. pallidoroseum*, *F. subglutinans*, *Penicillium* spp., *Rhizoctonia solani* and *Rhizopus* sp., were isolated and identified. Of the fungi isolated *Alternaria alternata*, *Aspergillus flavus*, *A. niger*, *Cephalosporium* sp., *Penicillium* spp., and *Rhizopus* sp., were found to be predominant. There does not appear to be any previous report of *Chaetomium globosum* and *Drechslera hawaiiensis* on wheat, whereas, *Absidia* sp., *Aspergillus sulphureus*, *Fusarium subglutinans* and *Rhizoctonia solani* appeared to be new report on wheat from Pakistan (Khanzada et al., 2002; Ahmed et al., 1992; Richardson, 1983; Ahmed et al., 1997; Ghaffar & Abbas, 1972; Glazek, 1997; Mirza & Qureshi, 1978; Weber et al., 2001).







Significant difference was observed in seed samples of wheat collected from Islamabad ( $p<0.001$ ) and Faisalabad ( $p<0.001$ ), whereas samples collected from Karachi showed insignificant difference. Methods for isolation of fungi showed significant difference in samples collected from Karachi ( $p<0.05$ ), Faisalabad ( $p<0.001$ ), Lahore ( $p<0.01$ ) and Islamabad ( $p<0.001$ ). Significant number ( $p<0.001$ ) of fungi were isolated from samples collected from all localities.

**b. Sorghum:** From 27 samples of sorghum, 14 genera and 23 species of fungi viz., *Alternaria alternata*, *Aspergillus* sp., *A. candidus*, *A. flavus*, *A. niger*, *A. sulphureus*, *Curvularia* sp., *C. lunata*, *Cladosporium* sp., *Drechslera* sp., *D. halodes*, *D. tetramera*, *D. hawaiiensis*, *Nigrospora oryzae*, *Trichoderma hamatum*, *Trichothecium roseum*, *Piptocephalis* sp., *Syncephalastrum racemosum*, *Fusarium moniliforme*, *F. subglutinans*, *Penicillium* spp., and *Rhizopus* sp. were isolated and identified. Of these, *Aspergillus candidus*, *A. flavus*, *A. niger*, *Penicillium* spp., *Piptocephalis* sp. and *Rhizopus* sp. were found to be predominant. There does not appear to be any previous report of *Aspergillus sulphureus*, *Nigrospora oryzae*, *Trichoderma hamatum*, *Fusarium subglutinans*, *Piptocephalis* sp., and *Syncephalastrum racemosum* on sorghum (Ahmed *et al.*, 1992; Ahmed *et al.*, 1997; Ghaffar & Abbas, 1972; Mirza & Qureshi, 1978; Williams and McDonald, 1983).

Significant difference was observed in seed samples of sorghum collected from Karachi ( $p<0.001$ ). Simple blotter and deep freezing methods for isolation of fungi showed significant difference in samples collected from Karachi ( $p<0.01$ ) and Lahore ( $p<0.01$ ), whereas nonsignificant difference was observed in samples collected from Islamabad and Faisalabad. Significant number ( $p<0.001$ ) fungi were isolated from samples collected from all localities except from Islamabad ( $p<0.01$ ).

**c. Barley:** From 14 samples of barley, 11 genera and 17 species of fungi viz., *Alternaria alternata*, *Aspergillus* sp., *A. candidus*, *A. flavus*, *A. niger*, *Cephalosporium* sp., *Cladosporium herbarum*, *Drechslera dematioidea*, *D. halodes*, *D. tetramera*, *Trichothecium roseum*, *Fusarium moniliforme*, *F. pallidoroseum*, *Penicillium* sp., *Stemphylium* sp., and *Ulocladium* sp. were isolated and identified. Of these *Alternaria alternata*, *Aspergillus niger*, *Penicillium* spp., *Cephalosporium* sp., and *Rhizopus* sp., were found to be predominant. *Aspergillus niger*, *Cephalosporium* sp., *Cladosporium herbarum*, *Drechslera dematioidea*, *D. tetramera*, *Trichothecium roseum*, *Stemphylium* sp., and *Ulocladium* sp., appeared to be new records on barley (Ahmed *et al.*, 1992; Ahmed *et al.*, 1997; Chong & Sheridan, 1982; Ghaffar & Abbas, 1972; Mirza & Qureshi, 1978).

Significant number ( $p<0.001$ ) of fungi were isolated from seed samples of barley collected from Karachi and Faisalabad. Significant difference ( $p<0.001$ ) was observed in simple blotter and deep freezing methods.

In the present work higher percentage of *Fusarium* spp., and *Drechslera* spp., was found where deep freezing method was used and mean infection range of *Fusarium* also increased (Fakhrunnisa & Hashmi, 1992). This may be partly due to the ease of observation and partly due to the fact that the dead embryo provides nourishment to the developing mycoflora. SNA medium appears satisfactory for the isolation and identification of *Fusarium* spp.

### References

Abdullah, S.K. and S.A. Kadhum. 1987. Seed mycoflora of *Sorghum bicolor* in Iraq. *Art Gulf J. Sci. Res.*, 5(3): 401-410.

Agarwal, V.K. and J.B. Sinclair. 1996. *Principles of Pathology*. 2<sup>nd</sup> edi, CRC Press, Inc., Boca Raton, Fl. 539 pp.

Ahmed S., S.H. Iqbal and A.N. Khalid. 1997. *Fungi of Pakistan*. Dept. Bot., University of the Punjab.

Ahmed, I., S. Iftikhar and A.R. Bhutta. 1992. Seed-borne microorganism in Pakistan Checklist 1991. PARC, Islamabad.

Anonymous. 1966. *International Rules for Seed Testing*, 1966. Proc. Int. Seed Test. Assoc., 31: 1-152.

Barnett, H.L. and B.B. Hunter. 1972. *Illustrated Genera of Imperfect Fungi*. Burgess Pub. Co., Minneapolis, Minnesota. 241pp.

Bateman, G.L. and H. Kwasna. 1999. Effects of number of winter wheat crops grown successively on fungal communities on wheat roots. *Applied Soil Ecology*, 13: 271-282.

Booth, C. 1971. *The Genus Fusarium*. Comonw. Mycol. Inst., Kew, Surrey, England. 237pp.

Chong, L.M. and J.E. Sheridan 1982. Mycoflora of barley (*Hordeum vulgare L.*) seed in New Zealand. *New Zealand Journal of Botany*, 20: 187-189.

Dawson, W.A.J.M. and G.L. Bateman. 2001. Fungal communities on roots of wheat and barley and effects of seed treatments containing fluquinconazole applied to control take-all. *Plant Pathology*, 50: 5-82.

Fakhrunnisa and M.H. Hashmi. 1992. Seed-borne mycoflora of corn, millet and paddy. pp. 125-129. In: Status of Plant Pathology in Pakistan. (Eds.): A. Ghaffar and S. Shahzad. Dept. Bot., Univ. Karachi, Karachi-75270, Pakistan.

Ghaffar, A. and S.Q. Abbas. 1972. Fungi of Pakistan Suppl. II. *Pak. J. Bot.*, 4: 195-208.

Glazek, M. 1997. Mycoflora of winter wheat seeds harvested from flooded commercial fields in South-Western Poland in 1997. Plant Protection Institute in Poznaniu, Sosnicowice Branch, Gliwicka St. 29:44-153 Sosnicowice, Poland.

Halt, M. 1994. *Aspergillus flavus* and aflatoxin B1 in flour production. *Eur. J. Epidermiol.*, 10(5): 555-8.

Hashmi, M.H. and U. Thrane. 1990. Mycotoxins and other secondary metabolites in species of *Fusarium* isolated from seeds of capsicum, coriander and fenugreek. *Pak. J. Bot.*, 22: 106-116.

Khanzada, K.A., M.A. Rajput, G.S. Shah, A.M. Lodhi and F. Mehboob. 2002. Effect of seed dressing fungicides for the control of seed borne mycoflora of Wheat. *Asian Journal of Plant Sciences*, 1(4): 441-444.

Kubiak K. and M. Korbas. 1999. Occurrence of fungal diseases on selected winter wheat cultivars. Postepy w Ochronie Roslin 39 (2), 801-804 [in Polish].

Limonard, T. 1968. Ecological aspect of seed health testing. *Proc. Int. Seed Test. Assoc.*, 33: 71-73.

Mirza, J.H. and M.S.A. Qureshi. 1978. *Fungi of Pakistan*. University of Agric., Faisalabad. Pakistan.

Nelson, P.E., T.A. Toussoun and W.F.O. Marasas. 1983. *Fusarium species. An Illustrated Manual of Identification*. The Pennsylvania State Univ. Press, Univ. Park, Pennsylvania. 203 pp.

Nirenberg, H. 1976. Untersuchungen über die morphologische und biologische Differenzierung in der *Fusarium* Sektion Liseola. Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft, Berlin-dehlena, 169: 1-117.

Nirenberg, H., H. Schmitz-Elsherif, C.I. Kling. 1994. Occurrence of Fusaria and some "blackening moulds" on durum wheat in Germany. 1. Incidence of *Fusarium* species. *Pflanzenkrankheiten und Pflanzenschutz*, 101: 449-459.

Richardson, M.J. 1983. *An annotated list of seed-borne diseases*, Supplement 2. Int. seed Test. Assoc., Zurich, Switzerland. 108pp.

Weber R., W. Kita, B. Hrynczuk, B. Runowska-Hrynczuk. 2001. Effects of tillage methods on the occurrence of culm base disease in several winter wheat cultivars. *Electronic J. Polish Agric. Univ., Agronomy*, 4(2):

Weber, R., B. Hrynczuk, B. Runowska-Hrynczuk and W. Kita. 2001. Influence of the mode of tillage on diseases of culm base in some winter wheat varieties, oats, and spring wheat. *J. Phytopathol.*, 149: 185-188.

Williams, R.J. and D. McDonald. 1983. Grain molds in the tropics: Problems and Importance. *Annual Review of Phytopathology*, 21: 153-178.

(Received for publication 7 March 2005)