

COMPARATIVE EFFECTS OF VARIOUS SALICYLIC ACID DERIVATIVES ON KEY GROWTH PARAMETERS AND SOME ENZYME ACTIVITIES IN SALINITY STRESSED MAIZE (*ZEA MAYS* L.) PLANTS

A. LEVENT TUNA¹, CENGİZ KAYA^{2*}, MURAT DIKİLİTAS³, İBRAHİM YOKAS⁴, BETÜL BURUN¹ AND HAKAN ALTUNLU⁴

¹Biology Department, Science and Arts Faculty, Mugla University, Mugla, Turkey

²Soil Science Department, Faculty of Agriculture, Harran University, Sanliurfa, Turkey

³Plant Protection Department, Faculty of Agriculture, Harran University, Sanliurfa, Turkey

⁴Technical Department, Ortaca Vocational School, Mugla University, Mugla, Turkey

*Corresponding author e-mail: c_kaya70@yahoo.com

Abstract

Salicylic acid, 5-Sulfo Salicylic acid and Acetylsalicylic acid are Salicylic acid derivatives. They differ in their substitution on the benzene ring and may have different effects on plant membranes. The effects of the derivatives of various Salicylic Acid [Salicylic acid (SA), 5-Sulfo Salicylic Acid (SSA) and Acetylsalicylic Acid (ASA)] on antioxidant enzyme activities, mineral element uptake, growth and some stress related parameters of maize (*Zea mays* L. cv. DK 684) plant grown in containers under salinity stress were investigated. Salicylic acid were applied by foliar treatments at five days interval. Treatments were: 1-) control, 2-) salt treatment 125 mM NaCl, 3-) 125 mM NaCl + 1 mM Salicylic Acid, 4-) 125 mM NaCl + 1 mM Sulfo Salicylic Acid, 5-) 125 mM NaCl + 1 mM Acetylsalicylic Acid, 6-) 125 mM NaCl + 2 mM Salicylic Acid, 7-) 125 mM NaCl + 2 mM Sulfo Salicylic Acid and 8-) 125 mM NaCl + 2 mM Acetylsalicylic Acid. Salt treatment reduced the plant growth, chlorophyll content, relative water content and ear of corn weight, but increased antioxidative enzymes and membrane permeability. Besides compared with the control group, nutrient uptakes of leaves and roots were inhibited by salt treatment. Tested parameters were generally positively affected by the applications of the salicylic acid derivatives compared to the salt treatment. For example, total chlorophyll, shoot dry matter, relative water content and ear of corn weight were ameliorated by 1 and 2 mM SSA, 1 mM ASA and 1 mM SSA treatments. The macro and micro element content of leaves and roots were generally increased by salicylic acid treatments compared to the salt treatment. Salicylic acid application seems to be more effective in the element uptakes than other ones. Salicylic acid treatments decreased antioxidant enzyme activities compared to the salt treatment. The data clearly shows that, the various derivatives of salicylic acid could protect maize plant from the detrimental effects of salt stress by improving physiological parameters tested such as relative water content, membrane permeability and nutrient status of plant.

Introduction

Salinity is one of the limiting environmental factors for crop production, and is becoming more serious issue as the intensity of agriculture increases. All around the world, about 100 million ha, or 5% of arable land has already been adversely affected by high salt concentrations which reduce crop growth and yield (Ghassemi *et al.*, 1995; Gunes *et al.*, 2007).

Salicylic acid (SA) (*o-hydroxybenzoic acid*), which belongs to a group of plant phenolics, is widely distributed in plants and is now considered as a hormone-like substance, which plays an important role in the regulation of plant growth and development (Raskin, 1992; Klessig & Malamy, 1994). During the last 20 years this

substance has drawn the attention of researchers because of its ability to induce systemic acquired resistance (SAR) in plants. At the present, considerable interest has been aroused by the ability of SA to produce a protective effect on plants under the action of stress factors of different abiotic nature. Thus considerable data have been obtained concerning the SA induced increase in the resistance of wheat seedlings to salinity (Shakirova & Bezrukova, 1997), and water deficit (Bezrukova *et al.*, 2001), of tomato and bean plants to low and high temperature (Senaratna *et al.*, 2000), as well as the injurious action of heavy metals on rice plants (Mishra & Choudhuri, 1999). Gomez *et al.*, (1993) and Rajasekaran & Blake (1999) reported a positive effect of SA on photosynthesis and plant growth under stress. Gomez *et al.*, (1993) observed greater economic yield of wheat cultivars grown under water stress when treated with SA. In maize plants, pretreatment with SA caused a decrease in net photosynthesis under normal growth conditions, but it activated some antioxidant enzymes (POX and GR), which in turn increased chilling tolerance in subsequent 2°C stress (Janda *et al.*, 2000). However, its exogenous application to the plants generates diverse physiological effects, such as inhibition of dry mass accumulation (Schettel & Balke, 1983), and control of ion uptake and their transport (Harper & Balke, 1981).

Salicylic acid (SA) has been reported to cause a multitude of effects on the morphology and physiology of plants (Pierpoint, 1994; Pancheva *et al.*, 1996) and to induce a protective mechanism enhancing resistance to biotic and abiotic stresses (Lopez-Delgado *et al.*, 1998). There is also evidence that SA can alter the antioxidant capacity in plants (Chen *et al.*, 1997; Rao *et al.*, 1997). Many studies support the SA-induced increases in the resistance of wheat and maize to salinity (Gunes *et al.*, 2007; Sakhabutdinova *et al.*, 2003; Shakirova & Bezrukova, 1997; Shakirova *et al.*, 2003) and osmotic stress (Bhupinder & Usha, 2003) and of rice on heavy metal stress (Mishra & Choudhuri, 1999).

In the present study, the effects of salicylic acid and its derivatives as a typical plant hormone, on the growth of the maize plant, nutrient uptake, antioxidant enzyme activities and some other stress related parameters were investigated.

Materials and Methods

Zea mays L. cv. DK 647, was used and the study was designed as three replicates according to trial plot in line with random plots design. Treatments were designed as follows: 1-) Control (only nutrient solution), 2-) NaCl 125 mM, 3-) NaCl 125 mM plus 1 mM SA, 4-) NaCl 125 mM plus 1 mM SSA, 5-) NaCl 125 mM plus 1 mM ASA, 6-) NaCl 125 mM plus 2 mM SA, 7-) NaCl 125 mM plus 2 mM SSA, 8-) NaCl 125 mM plus 2 mM ASA

Three seeds of maize were sown directly in plastic pots containing 8.0 kg of peat, perlite and sand mixture in equal ratios; following germination, plants were thinned to one plant per pot. The plants were fed with modified Hoagland-Arnon solution. The ratios of the minerals in this solution are as follows mg/l): 270 N, 30 P, 240 K, 200 Ca, 60 S, 50 Mg, 3 Fe, 0.5 Mn, 0.5 B, 0.02 Cu, 0.05 Zn. Until their height became 40cm, they had only been treated with water and nutritional substance and the nutritional solution was given to the control plants. When they became longer than 40 cm, Salicylic acid derivatives were sprayed to the leaves with five day-intervals. Each treatment was replicated three times and each replicate included 5 pots (i.e. 15 pots per treatment). The pH of the nutrient solution was adjusted to 6.5 with 0.1 mM KOH during the entire growing period. The volume of the nutrient solution applied to the root zone of the plants

ranged from 200 to 500 ml from the end of March to the middle of June each day depending on plant age. Plants were harvested 90 days after seedling emergence.

Chlorophyll determination: Prior to extraction, fresh leaf samples were cleaned with deionized water to remove any surface contamination. Chlorophyll extraction was carried out on fresh fully expanded leaf material; one g leaf sample was ground in 90% acetone using a pestle and mortar. The absorbance was measured with a UV/Visible spectrophotometer (Pye Unicam SP6-550, UK) and chlorophyll concentrations were calculated using the equation proposed by Strain & Svec (1966).

$$\text{Chl. a (mg ml}^{-1}\text{)} = 11.64 \times (\text{A663}) - 2.16 \times (\text{A645})$$

$$\text{Chl. b (mg ml}^{-1}\text{)} = 20.97 \times (\text{A645}) - 3.94 \times (\text{A663})$$

where (A663) and (A645) represent absorbance values read at 663 and 645 nm wavelengths, respectively.

Electrolyte leakage: This parameter was included in order to have more information on membrane stability and thereby on the relative ion content in the apoplastic space. Electrolyte leakage was assessed as described by Lutts *et al.*, (1996) using 9 young leaf discs for each treatment. Samples were washed three times with deionized water to remove surface-adhered electrolytes. Leaf discs were placed in closed vials containing 10 ml of deionized water and incubated at 25°C on a rotary shaker for 24 h; subsequently electrical conductivity of the solution (L_t) was determined. Samples were then autoclaved at 120°C for 20 min., and the last electrical conductivity (L_0) was obtained after equilibration at 25°C. The electrolyte leakage was defined as follows:

$$\text{Electrolyte leakage (\%)} = (L_t/L_0) \times 100$$

Proline determination: Proline was determined according to the method described by Bates *et al.*, (1973). Approximately 0.5 g of fresh leaf material was homogenized in 10 ml of 3% aqueous sulfosalicylic acid and filtered through Whatman's No. 2 filter paper. Two ml of the filtrate was mixed with 2 ml acid-ninhydrin and 2 ml of glacial acetic acid in a test tube. The mixture was placed in a water bath for 1 h at 100°C. The reaction mixture was extracted with 4 ml toluene and the chromophore containing toluene was aspirated, cooled to room temperature, and the absorbance was measured at 520 nm with a Shimadzu UV 1601 Spectrophotometer. Appropriate proline standards were included for the calculation of proline in the samples.

Leaf relative water content: Leaf relative water content (LRWC) was calculated based on the methods from Yamasaki & Dillenburg (1999). Two leaves of two randomly chosen plants per replicate were collected from the mid-sections of the plants in order to minimize age effects. Individual leaves were first removed from the stem and then weighed to obtain fresh mass (FM). In order to determine the turgid mass (TM), leaves were floated on distilled water inside a closed Petri dish. Maximum turgidity was determined by weighing leaves (after gently wiping the water from the leaf surface with tissue paper) until no further weight increase occurred. At the end of the imbibition period, leaf samples were placed in a pre-heated oven at 80°C for 48 h, in order to obtain dry mass (DM). All mass measurements were made using an analytical scale, with a

precision of 0.0001 g. Values of FM, TM and DM were used to calculate LRWC using the equation below:

$$\text{LRWC (\%)} = [(\text{FM-DM})/(\text{TM-DM})] \times 100$$

Protein content: Protein content in the enzyme extracts was determined according to Bradford (1976) using Bovine Serum Albumin V as a standard.

Enzyme Determination: Leaves (0.5 g) were homogenized in 50 mM sodium phosphate buffer (pH 7.0) containing 1% soluble polyvinyl pyrrolidine (PVP). The homogenate was centrifuged at 20,000 g for 15 min at 4°C and the supernatant used for assays of the activities of POX and SOD.

The activity of SOD was assayed by monitoring its ability to inhibit the photochemical reduction of NBT (Beauchamp and Fridovich, 1971). One unit of SOD was defined as the amount of enzyme necessary to inhibit the reduction of cytochrome c by 50%.

The activity of POD was assayed by adding aliquot of the tissue extract (100 µl) to 3 ml of assay solution, consisting of 3 ml of reaction mixture containing 13 mM guaiacol, 5 mM H₂O₂ and 50 mM Na-phosphate (pH 6.5) (Chance and Maehly, 1955). An increase of the optical density at 470 nm for 1 min at 25°C was recorded using a spectrophotometer. POD activity was expressed as change in absorbance min⁻¹ mg⁻¹ protein. The increase in A₄₇₀ was measured for 3 min and activity expressed as ΔA₄₇₀/mg protein/min.

Polyphenol oxidase (PPO) activity was assayed with 4-methylcatechol as a substrate according to the method of Zaubermann et al. (1991). Half gram of fresh leaf was ground with 10 ml of 0.1 mol/l sodium phosphate buffer (pH 6.8) and 0.2 g of polyvinylpyrrolidone (PVP, insoluble). After centrifugation at 19,000 g for 20 min, the supernatant was collected as the crude enzyme extract. The assay of the enzyme activity was performed using 1 ml of 0.1 mol/l sodium phosphate buffer (pH 6.8), 0.5 ml of 100 mmol/l 4-methylcatechol, and 0.5 ml enzyme solution. The increase in absorbance at 410 nm at 25 °C was recorded automatically for 5 min. One unit of enzyme activity was defined as an increase of 0.01 in absorbance per min per mg protein.

Statistics: Each pot was considered to be an experimental unit. Treatments were replicated three times and each replicate included 5 pots (i.e. 15 pots per treatment). The data for all parameters were statistically analyzed using the Statview-ANOVA test on computer. Statistically different groups were compared using an LSD test (*P*<0.05).

Results and Discussion

Key growth parameters: In the plants treated with salt, the amount of proline increased by %150 when compared to the control plants (Table 1). In NaCl treatment with the application of 1mM dose of SA, the increase in the proline continued; however, it decreased with 2mM SA dose. It is believed that 2mM dose of SA and its derivatives cause decrease in proline content by supporting the defense system of the plant. Proline indirectly causes increase in metabolic activation by providing osmoregulation for the plants under stress (Pesserakli & Huber, 1987). Proline amino acid has gained an

important position as a research subject in plant stress physiology. Although it is still under discussion, this compound has a defense mechanism supporting role for the plants under stress (Hare & Cress, 1997).

Relative water content (RWC) decreases with the application of salt compared to control treatment, but SA application increased it either moderately or remained stable (Table 1). This could be explained by transpiration decreasing effects of SA and its derivatives on leaves and epidermis (Aktaş, 2001). Agarwal *et al.*, (2005) reported that RWC capacity increased in wheat with SA treatment. It is known that salinity conditions decreases water uptake. For instance, in a study investigating plant-water relations in tomato plant and the ways of water uptake, it was found that when the tomato plant is watered with saltwater, a decrease both in its growth and its water uptake is observed (Romero-Aranda *et al.*, 2001).

Table 1. The effects of salicylic acid derivates on dry matter, proline and relative water content of maize plants grown at high NaCl.

Treatments	Shoot dry matter (%)	Root dry matter (%)	Proline μ moles/g FW	RWC %
Control	17.40a*	12.90a	7.44e	79.90a
NaCl	14.76b	11.76abc	18.35bc	67.46b
NaCl+ 1 mM SA	14.83b	11.33abc	20.25b	72.73ab
NaCl +1 mM SSA	14.03b	11.03c	22.85a	71.70ab
NaCl +1 mM ASA	15.60b	12.66ab	18.00c	68.36b
NaCl +2 mM SA	15.33b	11.23bc	11.77d	69.53b
NaCl +2 mM SSA	15.36b	11.96abc	11.07d	73.40ab
NaCl +2 mM ASA	15.30b	11.16bc	16.60c	72.10ab

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)

Table 2. The effects of salicylic acid derivates on stem height, stem diameter and ear of corn weight-number of maize grown at high NaCl.

Treatments	Stem height (cm)	Stem diameter (mm)	Ear of corn number (unit/plant)	Ear of corn weight (g/plant)
Control	230.8a*	18.433	1.88b	274.35a
NaCl	162.9c	16.313	2.44ab	158.13bc
NaCl+ 1 mM SA	164.5bc	17.443	2.77a	164.15bc
NaCl +1 mM SSA	159.8c	14.457	2.55a	184.51b
NaCl +1 mM ASA	165.8bc	16.923	2.55a	148.53c
NaCl +2 mM SA	168.8bc	18.467	2.66a	160.25cb
NaCl +2 mM SSA	179.3b	17.630	2.77a	169.61cb
NaCl +2 mM ASA	175.1bc	16.937	2.55a	160.02cb

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)

NaCl treatment reduced root and shoot dry matter. ASA treatment at 1 mM resulted in a moderate increase in root and shoot dry matter, but SA treatments did not result in significant change in this parameter (Table 1). The positive impact of SA on dry and fresh weight of the plant has been cited. For instance, 0.75 mM SA applied to wheat plant under stress increased both root and shoot length and dry and fresh weight. Moreover, SA given in the same dose, increased the yield in unsalted environment and had positive impacts on the physiological properties of the spike of the wheat plant. The chlorophyll a content of leaves under salinity conditions increased with 0.25 and 1 mM SA treatment (Arfan *et al.*, 2007).

In parallel with the vegetative growth deteriorations under stress, some physical parameters are also adversely affected (Blumwold, 2000). At the foremost of these come the stem height, stem diameter and productivity-related characteristics of the plant. Stem height and diameter were reduced by high salinity, but both stem height and stem diameter of the plant were positively affected by 2 mM dose of SA and its derivatives (Table 2). Another parameters adversely affected by salinity were ear of corn number and ear of corn weight. Salinity increased ear of corn number, but reduced ear of corn weight. Application of different most of chemical compounds used did not change those parameters except for 2mM SSA where slight increases were obtained. Supporting of rooting and speeding of vegetative growth by Salicylic acid and some compounds which are close analogues of it can explain the reason behind the positive impact seen on plant height and yield-related parameters as can be seen from Table 3 (Aktas, 2001; Agarwal *et al.*, 2005).

Salinity impaired membrane permeability increasing electrolyte leakage. However, application of salicylic acid (SA) partly maintained membrane permeability (Table 3). It has been noted that maintaining integrity of cellular membranes under stress conditions is considered an integral part of salinity tolerance mechanisms. Present study showed that SA reduced the amount of ion leakage in salt stressed maize plants indicating that SA treatment has facilitated the maintenance of membrane functions under stress conditions. Supporting evidence was shown when SA reduced electrolyte leakage salt stressed tomato leaves and in corn leaf, rice leaf, cucumber hypocotyl under chilling stress (Szltveit & Kang, 2001; Stevens *et al.*, 2006). Salinity reduced chlorophyll and carotenoid contents in maize plant compared to the control plants, but application of 1mM doses of SSA and ASA increased total chlorophyll when compared to salt group. The highest carotenoid content was reached in 1mM SSA dose (Table 3). There is a relation among chlorophyll content, photosynthesis activity and CO₂ fixation. The decreases seen in chlorophyll and carotenoid contents are the result of the indirect effect of NaCl. The decrease in net CO₂ fixation is the result of water shortage, closure of stomatas, accumulation of apoplast salt and mesophyll cells loosing turgor and direct toxicity of salt ions. All these negative situations are the indicators of the fact that general metabolic activity has been deteriorated (Pesserakli & Huber, 1987).

Enzyme activities: Salinity increased the concentrations of all antioxidant enzymes tested (SOD, PPO and POX), but in most cases, application of mentioned chemicals reduced those enzyme activities.

Among the investigated antioxidant enzymes, SOD increased with salt treatment but decreased with SA treatments (Table 4). Similar case is also true for PPO and POX. However, application of SA in 2 mM dose considerably increased POX activity different

effects of SA on anti-oxidant activity of plants have been reported. For example, in a study where the effects of SA on apoplastic enzyme activity in wheat plant leaves, it was found that with SA treatment, while apoplastic catalase enzyme activity decreased, apoplastic peroxidase and polyphenol oxidase activities increased (Tasgin *et al.*, 2006). Plants have antioxidants and antioxidative enzyme in varying amounts which protect (Asada & Takahashi, 1987; Ye *et al.*, 2000). Chloroplasts have antioxidative defense mechanism against toxic oxygen derivatives, and the foremost of these antioxidants are: vitamin E, vitamin C, glutathione, beta carotene and zeaxanthin. Enzymes such as superoxide dismutase (SOD), ascorbat peroxidase (APx), glutathione reductase (GR) and catalase (CAT) are known to be the most effective enzymes in the destruction of these radicals (Cakmak & Marschner, 1992; Cakmak, 1994; Gosset *et al.*, 1994; Dionisio-Sese & Tobita, 1998; Sreenivasulu *et al.*, 1999; Sreenivasulu *et al.*, 2000).

In another study, the effects of SA on oxidative stress and antioxidant enzyme activity in different wheat genotypes were investigated. SA, when compared to control plants, increased catalase, superoxide dismutase and ascorbat peroxidase enzyme activities. According to the authors; the reason behind the increase seen in antioxidant enzyme activities and decrease in stress is the increase in contents of chlorophyll, carotenoid and relative water, and this increased the total biomass of the plant (Agarwal *et al.*, 2005).

Nutrient content in leaf and root: It is known that SA and its derivatives may cause ion uptake thwarting effect in some plants. But the truth of this has not been tested in uncontrolled conditions. As complex bio-chemical events take place in the plant metabolism under stress, and as every plant has different degree of tolerance for stress, the mechanism of ion uptake may be under the impact of other factors, too. As can be seen in Table 5, 6, 7 and 8, separate macro and micro element levels were detected in the leaf and the root. According to the data provided in the literature, while usually in plants under NaCl stress macro element uptake and transportation are reduced, there is not sufficient information on micro element uptake and transportation. (Pancheva *et al.*, 1996; Pierpoint, 1994; Raskin, 1992). Lopez & Satti (1996) reported that the NaCl added to nutrition solution considerably decreased Ca and K contents in the leaves of tomato plants.

Table 3. The effects of Salicylic acid derivatives on membrane permeability (MP), chlorophyll and carotenoid contents in maize plants grown at high NaCl.

Treatments	MP %	Carotenoid (mg/g FW)	Chlorophyll a (mg/g FW)	Chlorophyll b (mg/g FW)	Total Chlorophyll (mg/g FW)
Control	29.80d*	2.54ab	1.64ab	0.580a	2.22ab
NaCl	50.23a	2.37ab	1.48ab	0.496ab	1.97b
NaCl+ 1 mM SA	39.40c	2.17b	1.56ab	0.510ab	2.07ab
NaCl +1 mM SSA	38.80c	2.78a	1.76a	0.583a	2.34a
NaCl +1 mM ASA	42.07bc	2.50ab	1.79a	0.590a	2.38a
NaCl +2 mM SA	44.97b	2.43ab	1.54ab	0.500ab	2.04b
NaCl +2 mM SSA	45.00b	2.34ab	1.54ab	0.486ab	2.02b
NaCl +2 mM ASA	38.70c	2.29b	1.36b	0.446b	1.81b

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)

Table 4. The effects of salicylic acid derivates on some antioxidant enzymes activities of maize grown at high NaCl.

Treatments	SOD enzyme activity (Unit/ mg Prot.)	PPO enzyme activity (Unit x100/mg prot.)	POX enzyme activity ($\Delta A_{470}/\text{min}/\text{mg prot.}$)
Control	30.32b	4.79b	47.84c
NaCl	38.10a	6.34a	61.15b
NaCl + 1 mM SA	35.71a	5.87b	52.38c
NaCl + 1 mM SSA	28.02b	5.26b	43.57d
NaCl + 1 mM ASA	27.05b	5.55b	60.68b
NaCl + 2 mM SA	15.36c	3.30c	115.59a
NaCl + 2 mM SSA	29.11b	5.60b	60.34b
NaCl + 2 mM ASA	26.30b	5.65b	41.21d

SOD: Superoxide dismutase, PPO: Polyphenol oxidase, POX: Peroxidase

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)**Table 5. The effects of salicylic acid derivates on macro elements (%) in the leaves of maize plants grown at high NaCl.**

Treatments	N	P	K	Ca	Mg	Na
Control	3.08a*	0.49	2.02b	1.14a	0.64a	0.018c
NaCl	2.47e	0.42	1.17f	0.54b	0.45c	0.27ab
NaCl + 1 mM SA	2.97b	0.49	1.92cd	0.65b	0.54abc	0.25b
NaCl + 1 mM SSA	2.82c	0.44	1.79e	0.60b	0.58ab	0.36a
NaCl + 1 mM ASA	2.97b	0.43	1.87d	0.55b	0.49bc	0.3ab
NaCl + 2 mM SA	2.97b	0.46	2.11a	0.62b	0.50bc	0.24b
NaCl + 2 mM SSA	2.63d	0.41	1.75e	0.55b	0.52abc	0.30ab
NaCl + 2 mM ASA	2.53de	0.44	1.97bc	0.51b	0.45c	0.32ab

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)**Table 6. The effects of salicylic acid derivates on micro elements (mg kg^{-1}) in the leaves of maize plants grown at high NaCl.**

Treatments	Cu	Fe	Mn	Zn
Control	16.5a	308a	162cd	170ab
NaCl	15.1b	221d	179b	154c
NaCl + 1 mM SA	13.6cd	229c	189a	162bc
NaCl + 1 mM SSA	14.1c	233b	195a	172ab
NaCl + 1 mM ASA	14.2bc	229c	144e	173ab
NaCl + 2 mM SA	13.9cd	228c	158d	173ab
NaCl + 2 mM SSA	13.4cd	236b	189a	177a
NaCl + 2 mM ASA	13.1d	229c	165c	175ab

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)

Table 7. The effects of salicylic acid derivates on macro elements in the roots of maize plants grown at high NaCl.

Treatments	N	P	K	Ca	Mg	Na
Control	1.35a*	0.55a	3.24a	1.73a	2.11a	2.17g
NaCl	1.17b	0.17b	0.32bc	0.35e	0.42d	4.76a
NaCl + 1 mM SA	1.19ab	0.22b	0.37b	0.69c	0.59c	3.15d
NaCl + 1 mM SSA	1.24ab	0.20b	0.35b	0.57cd	0.43d	4.14b
NaCl + 1 mM ASA	1.19ab	0.16b	0.21de	0.49d	0.43d	3.52c
NaCl + 2 mM SA	1.20ab	0.22b	0.28bcd	0.92b	0.88b	2.54f
NaCl + 2 mM SSA	1.21ab	0.16b	0.18e	0.58cd	0.82b	3.21d
NaCl + 2 mM ASA	1.11b	0.19b	0.23cde	0.58cd	0.48d	2.84e

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)**Table 8. The effects of salicylic acid derivates on the micro elements in the roots of maize plants grown at high NaCl.**

Treatments	Cu	Fe	Mn	Zn
Control	28.2a	505c	45abc	160a
NaCl	15.6b	263e	33bcd	28b
NaCl + 1 mM SA	10.4b	191g	50a	31b
NaCl + 1 mM SSA	12.5b	550b	44abcd	19b
NaCl + 1 mM ASA	10.5b	249f	39abcd	19b
NaCl + 2 mM SA	15.9b	474d	48ab	29b
NaCl + 2 mM SSA	27.0a	640a	30cd	23b
NaCl + 2 mM ASA	33.0a	190g	28d	28b

SA: Salicylic Acid, SSA: 5-sulfo Salicylic Acid, ASA: Acetylsalicylic Acid, EC: Electrical conductivity

*Within each column, same letter indicates no significant difference between treatments ($P<0.05$)

As can be understood from the investigation of the tables, while SA and its derivatives given in addition to NaCl remain below the control, they are generally influential on micro element contents of the leaves and roots. This situation is more evident in the leaf in terms of N and K and in the root in terms of Ca and Mg. Among the SA derivatives, the one best competing with Na is Salicylic acid. When micro elements are looked into, while the best reaction in the leaf is given by Zn, the other micro elements reveals a fluctuating pattern in both the leaf and root.

There are studies showing that Salicylic acid improves the quality in plants. There is an indirect relation between the quality and nourishment. In a study conducted on *Prunus persica* (L.) it was found that with 1 mM SA treatment, the destruction caused by cold decreases and rotting index and fruit hardness are improved. As known, especially in fruits, Ca element is responsible for cellular stabilization and quality. In this study SA served a function similar to that of Ca (Wang *et al.*, 2006).

In another study, the effects of SA treatments ranging from 0.1 to 1 mM on the plant's element uptake, its growth and membrane permeability in maize plant under salinity stress were investigated. With 1 mM SA treatment, the membrane permeability considerably decreased when compared to NaCl group. While with SA treatment, Na and Cl uptake of the plant was strongly inhibited, N, Mg, Fe, Mn and Cu uptake was considerably stimulated. According to the researchers, SA took the role of an important hormone fostering the resistance against stress in the plant (Gunes *et al.*, in press).

In few studies, it has been reported that SA exhibits a synergisms with Ca in the plant cell. In a study conducted on grape plant, under normal circumstances, while the electrolyte leakage ratio decreased, cytosolic contents were found to be higher compared to the control. According to the researchers' statements, inner-cell Ca balance (homeostasis) and antioxidant system improves with the exogenous application of SA in plants under normal circumstances or under stress (Wang & Li, 2006).

Various physiological and biochemical effects of SA on plant systems have been well documented (Raskin, 1992; Cameron, 2000). However, studies related to ion concentration and uptake are relatively lacking. Aly & Soliman (1998) studied the effect of SA on iron uptake in soybean genotypes. They found that SA was effective in correcting iron chlorosis in soybean genotypes grown in calcareous soils. Al-Hakimi & Hamada (2001) also observed similar effects of SA in the Na, K, Ca and Mg content of wheat plants grown under salinity. Positive effects of SA on the ion uptake and inhibitory effects on Na and Cl uptake should also be responsible for managing salinity of maize plants (Gunes *et al.*, in press).

Conclusions

In the maize plant under the salinity stress, growth-related properties were adversely affected and anti-oxidant enzyme activities increased. SA derivatives given through the leaf, especially in lower doses, positively affected the growth parameters when compared to the control plants. Macro and micro element uptake of the maize plant generally increased by application of SA derivatives compared to salinity-stressed plants. Furthermore, SA treatments generally reduced antioxidant enzyme activities compared to plants at high NaCl. There seems to be no significant difference with regards to their effects on the plant among salicylic acid derivatives.

The obtained data reveal that salicylic acid derivatives which are found in the plants very commonly, and having hormone-like effect, can decrease the adverse effects of salinity stress, especially in low dose applications.

Acknowledgements

Authors wish to thank Mugla and Harran University, Scientific Research Projects Committee for supporting this research.

References

- Agarwal, S., R.K. Sairam, G.C. Srivastava and R.C. Meena. 2005. Changes in antioxidant enzymes activity and oxidative stres by abscisic acid and salicylic acid in wheat genotypes. *Biologia Plant.*, 49: 541-550.
- Aktaş, Y.L. 2001. *Vitis vinifera'da Salisilik asit uygulamasının yaprak proteinleri içeriği üzerine etkileri*. EÜ Fen Bil Enst Biyoloji ABD Doktora Tezi, İzmir
- Al-Hakimi, A.M.A. and A.M. Hamada. 2001. Counteraction of salinity stress on wheat plants by grain soaking in ascorbic acid, thiamin or sodium salicylate. *Biol Plant.*, 44: 253-61.
- Aly, S.S.M. and S.M. Soliman. 1998. Impact of some organic acids on correcting iron chlorosis in two soybean genotypes grown in calcareous soil. *Nutr Cycling Agroecosyst*, 51: 185-91.
- Arfan, M., H.R. Athar and M. Ashraf. 2007. Does exogenous application of salicylic acid through the rooting medium modulate growth and photosynthetic capacity in two differently adapted spring wheat cultivars under salt stress? *Journal of Plant Physiology*; (In press).

Asada, K and M Takahashi. 1987. Production and scavenging of active oxygen radicals in photosynthesis. In: *Photoinhibition*. (Eds.): DJ Kyle *et al.*, Elsevier, Amsterdam, 227-297.

Beauchamp, C. and Fridovich, I. 1971. Superoxide dismutase improved assays and an assay applicable to acrylamide gels, *Anal. Biochem.* 444: 276-287

Bezrukova, M.V., R. Sakhabutdinova, R.A. Fatkhutdinova, I. Kyldiarova and F. Shakirova. 2001. The role of hormonal changes in protective action of salicylic acid on growth of wheat seedlings under water deficit. *Agrochemiya (Russ)*, 2:51-54.

Bhupinder, S. and K. Usha. 2003. Salicylic acid induced physiological and biochemical changes in wheat seedlings under water stress. *Plant Growth Regul.*, 39: 137-41.

Blumwold, E. 2000. Sodium transport and salt tolerance in plants. *Current Opinion in Cell Biology*, 12: 431-434.

Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal Biochem*, 72: 248-254.

Cakmak, I. 1994. Activity of ascorbate-dependent H₂O₂-scavenging enzymes and leaf chlorosis are enhanced in magnesium -and potassium- deficient leaves, but not in phosphorus-deficient leaves. *J. Exp. Bot.*, 45: 1259-1266.

Cakmak, I. and H. Marschner. 1992. Magnesium deficiency and high-light intensity enhance activities of superoxide dismutase, ascorbate peroxidase and glutathione reductase in bean leaves. *Plant Physiol.*, 98: 1222-1227.

Cameron, R.K. 2000. Salicylic acid and its role in plant defense responses: what do we really know? *Physiol. Mol. Plant. Pathol.*, 56: 9-13.

Chance, B. and Maehly, C. 1955. Assay of catalase and peroxidases, *Methods Enzymol.* 11: 764-775

Chen, Z., S. Iyer, A. Caplan, D.F. Klessig and B. Fan. 1997. Differential accumulation of salicylic acid and salicylic acid-sensitive catalase in different rice tissues. *Plant Physiol.*, 114: 193-201.

Dionisio-Sese, M.L. and S. Tobita. 1998. Antioxidant responses of rice seedlings to salinity stress. *J. Plant Sci.*, 135: 1-9.

Ghassemi, F., A.J. Jakeman and H.A. Nix. 1995. *Salinisation of land and water resources*. Wallingford, UK: CAB International.

Gomez, L., L. Blanca and C.S. Antonio. 1993. Evidence of the beneficent action of the Acetylsalicylic acid on wheat genotypes yield under restricted irrigation. In: *Proc. Scientific Meeting on Forestry, Livestock and Agriculture Mexico*, p. 112.

Gosset, D.R., E.P., Millhollon and C. Lucas. 1994. Antioxidant response to NaCl stress in salt-tolerant and salt-sensitive cultivars of cotton. *Crop Sci.*, 34: 706-714.

Gunes, A., A. Inal, M. Alpaslan, F. Eraslan, E.G. Bagci and N. Cicek. 2007. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays L.*) grown under salinity. *Journal of Plant Physiology*; (In press).

Hare, P.D. and W.A. Cress. 1997. Metabolic implications of stress-induced proline accumulation in plants. *Plant Growth Regulation*, 21: 79-102.

Harper, J.R and N.E. Balke. 1981. Characterization of the inhibition of K⁺ absorption in oats roots by salicylic acid. *Plant Physiol.*, 68: 1349-1353.

Janda, T., G. Szalai Antunovics, Z.E. Horvath and E. Paldi. 2000. Effect of benzoic acid and aspirin on chilling tolerance and photosynthesis in young maize plants. *Maydica*, 45: 29-33.

Klessig, D.F. and J. Malamy. 1994. The salicylic acid signal in plants. *Plant Mol Biol.*, 26: 1439-1458.

Lopez, M.V. and S.M.E. Satti. 1996. Calcium and potassium enhanced growth and yield of tomato under sodium chloride stress. *Plant Sci.*, 114: 19-27.

Lopez-Delgado, H., J.F. Dat, H. Foyer and I.M. Scott. 1998. Induction of thermotolerance in potato microplants by acetylsalicylic acid. *J. Exp. Bot.*, 49: 713-720.

Mishra, A. and M.A. Choudhuri. 1999. Effect of salicylic acid on heavy metal-induced membrane deterioration mediated by lipoxygenase in rice. *Biol. Plant.*, 42: 409-415.

Pancheva, T.V., L.P. Popova and A.N. Uzunova. 1996. Effect of salicylic acid on growth and photosynthesis in barley plants. *J. Plant Physiol.*, 149: 57-63.

Pessarakli, M. and J.T. Huber. 1987. Dry matter, yield, nitrogen absorption, and water uptake by sweet corn under salt stress. *J. Plant Nutr.*, 12: 279-290.

Pierpoint, W.S. 1994. Salicylic acid and its derivatives in plants: medicines, metabolites and messenger molecules. *Bot. Res.*, 20: 163-235.

Rajasekaran, LR and T.J. Blake. 1999. New plant growth regulators protect photosynthesis and enhance growth under drought of jack pine seedlings. *J. Plant Growth Reg.*, 18: 175-181.

Rao, M.V., G. Paliyath, P. Ormrod, D.P. Murr and C.B. Watkins. 1997. Influence of salicylic acid on H₂O₂ production, oxidative stress, and H₂O₂-metabolizing enzymes. *Plant Physiol.*, 115: 137-149.

Raskin, I. 1992. Role of salicylic acid in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 43: 439-463.

Romero-Aranda, R., T. Soria and J. Cuartero. 2001. Tomato plant water uptake and plant water relationships under saline growth conditions. *Plant Sci.*, 160: 265-272.

Sakhabutdinova, A.R., D.R. Fatkhutdinova, M.V. Bezrukova and F.M. Shakirova. 2003. Salicylic acid prevents damaging action of stress factors on wheat plants. *Bulg. J. Plant Physiol.*, 214-219. [special issue].

Schettel, N.L. and N.E. Balke. 1983. Plant growth response to several allelopathic chemicals. *Weed Sci.*, 31: 293-298.

Senaratna, T., D. Touchell, E. Bunn and K. Dixon. 2000. Acetylsalicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plant. *Plant Growth Reg.*, 30: 157-161.

Shakirova, F.M. and M.V. Bezrukova. 1997. Induction of wheat resistance against environmental salinization by salicylic acid. *Biol., Bull.*, 24: 109-112.

Shakirova, F.M., A.R. Sakhabutdinova, M.V. Bezrukova, R.A. Fathutdinova and D.R. Fathutdinova. 2003. Changes in hormonal status of wheat seedlings induced by salicylic acid and salinity. *Plant Sci.*, 164: 317-22.

Sreenivasulu, N., B. Grimm, U. Wobus and W. Weschke. 2000. Differential response of antioxidant compounds to salinity stress in salt-tolerant and salt-sensitive seedlings of fox-tail millet (*Setaria italica*). *Physiol. Plant.*, 109: 435-442.

Sreenivasulu, N., S. Ramanjulu, K. Ramachandra-Kini, H.S. Prakash, H. Shekar-Shetti, H.S. Savithri and C. Sudhakar. 1999. Total peroxidase activity and peroxidase isoforms as modified by salt stress in two cultivars of fox-tail millet with differential salt tolerance. *Plant Sci.*, 141: 1-9.

Stevens, J., T. Senaratna and K. Sivasithamparam. 2006. Salicylic acid induces salinity tolerance in tomato (*Lycopersicon esculentum* cv. Roma): associated changes in gas exchange, water relations and membrane stabilisation. *Plant Growth Regul.*, 49: 77-83.

Szltveit, M.E. and H. Kang. 2001. Activity of enzymatic antioxidant defense systems in chilled and heat shocked cucumber seedling radicles. *Physiol. Plant.*, 113: 548-556.

Tasgin, E., O. Atici, B. Nalbantoglu and L.P. Popova. 2006. Effects of salicylic acid and cold treatments on protein levels and on the activities of antioxidant enzymes in the apoplast of winter wheat leaves. *Phytochemistry*, 67: 710-715.

Wang, L., S. Chena, W. Kong, S. Li and D.D. Archbold. 2006. Salicylic acid pretreatment alleviates chilling injury and affects the antioxidant system and heat shock proteins of peaches during cold storage. *Postharvest Biology and Technology*, 41: 244-251.

Wang, L.J. and S.H. Li. 2006. Salicylic acid-induced heat or cold tolerance in relation to Ca²⁺ homeostasis and antioxidant systems in young grape plants. *Plant Sci.*, 170: 685-694.

Ye, Z., R. Rodriguez, A. Tran, H. Hoang, D.D. Los Santos and S. Brown. 2000. The developmental transition to flowering repesesses ascorbate peroxidase activity and induced enzymatic lipid peroxidation in leaf tissue in *Arabidopsis thaliana*. *Plant Sci.*, 158: 115-127.

Zauberman, G. Ronen, R. Akerman, M. Weksler, A. Rot, I. and Fuchs, Y. 1991. Postharvest retention of the red color of litchi fruit pericarp, *Sci. Hort.*, 47: 89-97.

(Received for publication 2 February 2007)