

## MEDICINAL PLANTS OF USHERAI VALLEY, DIR, NWFP, PAKISTAN

ALI HAZRAT\*, JEHANDAR SHAH, SHUJAAT AHMAD, MOHAMMAD NISAR,  
ABDUL KHALIQ JAN AND SIKANDAR

*Department of Botany and Research Centre,  
Shaheed Benazir Bhutto University and University of Malakand, Malakand, Pakistan.*

### Abstract

This research is based on the results of an ethno-botanical research conducted in Usherai Valley. The main objective was to enlist the wealth of medicinal plants. In total 50 species, belonging to 32 families of wild herbs, shrubs and trees were found to be used as medicinal plants by the inhabitants in the valley.

### Introduction

Usherai Valley is situated in the east side of Dir. It is 70km away from Timergara on the main Dir Chitral road. The area is fertile for agriculture. The total area covered by this Valley is 3992.67 hectares. The major part of the Usherai Valley is hilly. The whole area is generally covered with forests. The climate is extremely cold in winter and slightly warm in summer. The mean maximum and minimum temperature in the month of January has been recorded as 6.22°C and -2.39°C respectively (Ghafoor, 2002). The research area is rich with medicinal plants and the local people of the area widely use them as a remedy for various ailments. This remote area was selected to revive the old tradition because in past there was deficiency of doctors and the Hakims resorted to different medicinal plants as a treatment to different diseases. (Arshad, 1999; Shinwari, 2002-2003).

The eastern medicine practiced in Pakistan comprises of three systems Chinese, Ayurvedic and Greco Arabic. The recipes of medicines used in these systems are derived from both organic and inorganic sources. The knowledge of drugs goes back to prehistoric times. Records of ancient civilizations show that a considerable number of drugs, used by modern doctors, were already in use of Egyptians, Babylonians, Greeks, Romans, Chinese and people of the subcontinent (India and Pakistan). (Khan, 1991).

### Materials and Methods

Regular study trips were made to representative areas and collections were completed in their flowering season from March 2006- 2007. During these trips, different plants were collected, dried, documented and were identified both by comparing them with herbarium specimen and with the help of Flora of Pakistan. (Stewart, 1967, 1982). Specimen sets were given to the Herbarium of Islamia College University Peshawar and herbarium of Shaheed Benazir Bhutto University (one each) for future reference. Through a questionnaire, medicinal plants usage data was collected from local people and Hakims that practice the medicine regularly.

\*E-mail: aligom@gmail.com, ali\_hazrat8@yahoo.com, botanist2005@yahoo.com





## Results and Discussion

The research revealed that local folk utilize 50 species of plants, belonging to 32 families for various purposes (Table 1). The people of the valley are generally ignorant about the medicinal and economic importance of these plants. Out of the 50 medicinal plants, only 30 species were known to the local people and the rest of the species of high medicinal and economic values were completely unknown to the local community of the area. (Ahmad, 2000; Rehman and Ghafoor, 2002).

## References

Ahmad, H. 2000. The medicinal plants of Tharparker, WWF – Peshawar, Pakistan.

Arshad, M. and S. Akram. 1999. Medicinal plants of University of Arid Agriculture, Rawalpindi. *Hamdard Med.*, 42:46-49.

Choudhary, M., S. Ahmad, A. Ali, H. Sher and S. Malik. 2000. Technical Report: Market study of medicinal herbs in Malakand, Peshawar, Lahore and Karachi. SDC-Inter co-operation, Peshawar.

Ghufran, m. A., r. A. Qureshi, a. Batool, j. Guilford, l. E. Marlar, t. P. Kondratyuk, l. C. Chang, and j. M. Pezzuto. 2008. Evaluation of selected indigenous medicinal plants from the western himalayas for cytotoxicity and as potential cancer chemopreventive agents. *Pharmaceutical biology* (accepted)

Khan, a.a. 1991. Marketing of herbal drugs and its problems, proceedings of the national workshop on appropriate uses of medicinal plants in traditional medicines. Nih, islamabad. 69-72.

Qureshi, r. A., m. A. Ghufran, s. A. Gilani, z. Yousaf, g. Abbas and a. Batool. 2009. Indigenous medicinal plants used by local women in southern himalayan region of pakistan. *Pakistan journal of botany*, 41(1): 19-25.

Rehman, m. And s. Ghafoor. 2002. The natural resources and plant ecology of mount elum district swat. Consultancy report. Wwf-p, peshawar.

Shaheen F., M. Ahmad, M.T. Khan, S. Jalil, A. Ejaz, M.N. Sultankhodjaev, M. Arfan, M.I. Choudhary and Atta-ur-Rahman. 2005. Alkaloids of *Aconitum laeve* and their anti-inflammatory antioxidant and tyrosinase inhibition activities. *Phytochemistry*, Apr; 66(8): 935-40.

Subedar Pandey, Rekha Kushwaha, Om Prakas, Amita Bhattacharya and P.S. Ahuja. 2007. Ex situ conservation of *Aconitum heterophyllum* Wall.—an endangered medicinal plant of the Himalaya through mass propagation and its effect on growth and alkaloid content. Division of Biotechnology, Institute of Himalayan Bioresource Technology, Palampur-176061, H. P., India. 3: 127-135 Cambridge University Press doi: 10.1079/PGR200570.

Shinwari, Z.K. and S.S. Gilani 2003. Sustainable Haverst of medicinal plants, Bulashbar Nallah, Astor, Gilgit. *J. Ethno Pharmacology*, 84: 289-298.

Shinwari, Z.K., S.S. Gilani and M. Akhlas. 2002. Sustainable harvest of medicinal plants at Bar and Shinakii valleys, Gilgit. Consultancy Report: WWF-P.

Stewart, R.R. 1967. Checklist of plants Swat state, Northwest Pakistan. *Pak. J. For.*, 4(2): 457-528.

Stewart, R.R. 1982. History and exploration of plants in Pakistan and adjoining areas, National Herbarium, NARC, Islamabad.

Sudhir .K. Uniyal and O. P. Sati. 2001. Triterpenoid saponins from roots of *Clematis grata*. *Phytochemistry*, Volume 31, Issue 4, Pages 1427-1428.

(Received for publication 9 February 2009)